

ISSN 2536-4898
Volume 29
Issue 2
June 2019



Turkish Journal of **COLORECTAL DISEASE**

Official Journal of the Turkish Society of Colon and Rectal Surgery

Turkish Journal of COLORECTAL DISEASE



**Owner on Behalf of the Turkish Society of Colon and Rectal Surgery
/Türk Kolon ve Rektum Cerrahi Derneği Adına İmtiyaz Sahibi**
Selman SÖKMEN, M.D.

Dokuz Eylül University Faculty of Medicine, Department of General Surgery, İzmir, Turkey

Responsible Manager/Sorumlu Yazı İşleri
Tahsin ÇOLAK, M.D.

Mersin University Faculty of Medicine, Department of General Surgery, Mersin, Turkey

Editor-in-Chief/Baş Editör
Tahsin Çolak, M.D.

Mersin University Faculty of Medicine, Department of General Surgery, Mersin, Turkey
E-mail: colaktahsin@yahoo.com ORCID ID: orcid.org/0000-0002-7253-5608

Associate Editors/Editör Yardımcıları
Ahmet Rencüzoğulları, M.D.

Çukurova University Faculty of Medicine, Department of General Surgery, Adana, Turkey
ORCID ID: orcid.org/0000-0002-5993-9536

Ahmet Ziya Balta, M.D.
Haydarpaşa Sultan Abdülhamit Han Training and Research Hospital, İstanbul, Turkey
ORCID ID: orcid.org/0000-0002-4143-4169

İlker Sücüllü, M.D.
Haydarpaşa Sultan Abdülhamit Han Training and Research Hospital, İstanbul, Turkey
E-mail: ilkersucullu@gmail.com ORCID ID: orcid.org/0000-0002-5285-3051

Sezai Leventoğlu, M.D.
Gazi University Faculty of Medicine, Department of General Surgery, Ankara, Turkey
ORCID ID: orcid.org/0000-0003-0680-0589

Past Editors/Geçmiş Editörler

Erman Aytaç, M.D.
Ersin Öztürk, M.D., PhD.
Rasim Gençosmanoğlu, M.D.
Sezai Demirbaş, M.D.
Uğur Sungurtekin, M.D.
Fatma Ayça Gültekin, M.D.
Hüseyin Sinan, M.D.
M. Özgür Türkmenoğlu, M.D.
B. Bülent Menteş, M.D.
Kemal Alemdaroğlu, M.D.

Statistic Editor/İstatistik Danışmanı
Emine Arzu Okul, PhD.

Language Editor/Dil Editörü
Bedia Marangozoğlu, İstanbul, Turkey

Translator/Çevirmen
Bedia Marangozoğlu, İstanbul, Turkey

All inquiries should be addressed to

TURKISH JOURNAL OF COLORECTAL DISEASE

Address: Latilokum Sk. Alphan İşhanı No: 3 Kat: 2 Mecidiyeköy Şişli, İstanbul, Turkey
Phone: +90 212 356 01 75-76-77 Gsm: +90 532 300 72 36 Fax: +90 212 356 01 78

Online Manuscript: www.journalagent.com/krhd Web page: www.turkishjcrd.com E-mail: info@turkishjcrd.com

∞ All rights are reserved. Rights to the use and reproduction, including in the electronic media, of all communications, papers, photographs and illustrations appearing in this journal belong to the Turkish Journal of Colorectal Disease. Reproduction without prior written permission of part or all of any material is forbidden. The journal complies with the Professional Principles of the Press. The paper used to print this journal conforms to ISO 9706: 1994 standard (Requirements for Permanence). The National Library of Medicine suggests that biomedical publications be printed on acid-free paper (alkaline paper).

Reviewing the articles' conformity to the publishing standards of the Journal, typesetting, reviewing and editing the manuscripts and abstracts in English and publishing process are realized by Galenos.



Galenos Publishing House
Owner and Publisher
Erkan Mor

Publication Coordinator
Burak Sever

Web Coordinators
Turgay Akpınar

Graphics Department
Ayda Alaca
Çiğdem Birinci
Gülşah Özgül

Finance Coordinator
Sevinç Çakmak

Project Coordinators
Eda Kolukisa
Esra Semerci
Hatice Balta
Zeynep Altındağ

Project Assistants
Duygu Yıldırım
Gamze Aksoy
Melike Eren
Nurcan Acarçay
Pelin Bulut
Saliha Tuğçe Güdücü

Research&Development
Kerim Sancar Ölmez
Mert Can Köse

Publisher Contact

Address: Molla Gürani Mah. Kaçamak Sk.
No: 21/1 34093 İstanbul, Turkey
Phone: +90 (212) 621 99 25 Fax: +90 (212) 621 99 27
E-mail: info@galenos.com.tr/yayin@galenos.com.tr
Web: www.galenos.com.tr **Publisher Certificate Number: 14521**

Printing at: Üniform Basım San. ve Turizm Ltd. Şti.
Matbaacılar Sanayi Sitesi 1. Cad. No: 114
34204 Bağcılar, İstanbul, Turkey
Phone: +90 (212) 429 10 00 **Certificate Number: 42419**

Printing Date: June 2019
ISSN: 2536-4898 E-ISSN: 2536-4901
International scientific journal published quarterly.

Turkish Journal of COLORECTAL DISEASE



Reviewer Board

- Abdullah Zorluoğlu, M.D. (Acıbadem University Faculty of Medicine, Bursa)
Acar Aren, M.D. (İstanbul Training and Research Hospital, İstanbul)
Adil Baykan, M.D. (Medistate Hospital, Clinic of General Surgery, İstanbul)
Ahmet Özbal, M.D. (İstanbul University Cerrahpaşa Faculty of Medicine, İstanbul)
Alper Sözütek, M.D. (Numune Training and Research Hospital, Adana)
Andre D'Hoore, M.D. (University Hospitals Leuven, Belgium)
Andres Mellgren, M.D. (University of Illinois College of Medicine, Chicago, USA)
Angelita Habr - Gama, M.D. (University of Sao Paulo School of Medicine, Sao Paulo, Brazil)
Ann C. Lowry, M.D. (University of Minnesota, Minneapolis, USA)
Ayhan Kuzu, M.D. (Ankara University Faculty of Medicine, Ankara)
Bilgi Baca, M.D. (Acıbadem University Faculty of Medicine, İstanbul)
Bırol Bostancı, M.D. (Yüksek İhtisas Training and Research Hospital, Ankara)
Bülent Erkek, M.D. (Ankara University Faculty of Medicine, Ankara)
Bülent Menteş, M.D. (Memorial Hospital, Ankara)
Cem Kaan Parsak, M.D. (Çukurova University Faculty of Medicine, Adana)
Cem Terzi, M.D. (Dokuz Eylül University Faculty of Medicine, İzmir)
Cemalettin Ertekin, M.D. (İstanbul University Faculty of Medicine, İstanbul)
Cemil Çalışkan, M.D. (Ege University Faculty of Medicine, İzmir)
Cihangir Akyol, M.D. (Ankara University Faculty of Medicine, Ankara)
Cüneyt Kayaalp, M.D. (İnönü University Faculty of Medicine, Malatya)
Durkaya Ören, M.D. (Atatürk University Faculty of Medicine, Erzurum)
Dursun Buğra, M.D. (Koç University Faculty of Medicine, İstanbul)
Ediz Altunlu, M.D. (Florence Nightingale Hospital, İstanbul)
Emre Balık, M.D. (Koç University Faculty of Medicine, İstanbul)
Emre Canda, M.D. (Dokuz Eylül University Faculty of Medicine, İzmir)
Emre Görgün, M.D. (Cleveland Clinic, Ohio, USA)
Enis Yüney, M.D. (Okmeydanı Training and Research Hospital, İstanbul)
Eray Kara, M.D. (Celal Bayar University Faculty of Medicine, İstanbul)
Erdogan Sözüer, M.D. (Erciyes University Faculty of Medicine, Kayseri)
Ergün Yücel, M.D. (Haydarpaşa Sultan Abdülhamid Han Training and Research Hospital, İstanbul)
Erhun Eyüboğlu, M.D. (Kemerburgaz University Faculty of Medicine, İstanbul)
Erman Aytac, M.D. (Acıbadem University Faculty of Medicine, İstanbul)
Ersin Öztürk, M.D. (Medicana Hospital, Bursa)
Ethem Geçim, M.D. (Ankara University Faculty of Medicine, Ankara)
Faramarz Pakravan, M.D. (Center of Coloproctology, Duesseldorf, Germany)
Feza Karakayalı, M.D. (Başkent University Faculty of Medicine, İstanbul)
Feza Remzi, M.D. (New York University, Langone Medical Center, New York, USA)
Gökhan Yağcı, M.D. (Medicana Hospitals, Ankara)
Haldun Gündoğdu, M.D. (Atatürk Training and Research Hospital, Ankara)
Halis Dokgöz, M.D. (Mersin University Faculty of Medicine, Mersin)
Hiroki Ohge, M.D. (Hiroshima University Hospital, Hiroshima, Japan)
Hovsep Hazar, M.D. (Marmara University Faculty of Medicine, İstanbul)
İlyas Başkonuş, M.D. (Gaziantep University Faculty of Medicine, Gaziantep)
İsmail Cem Eray, M.D. (Çukurova University Faculty of Medicine, Adana)
İsmail Hamzaoğlu, M.D. (Acıbadem University Faculty of Medicine, İstanbul)
Janindra Warusavitarn, M.D. (St Mark's Hospital and Academic Institute, Harrow, United Kingdom)
Julio Garcia-Aguilar, M.D. (Memorial Sloan Kettering Cancer Center, New York, USA)
Khaled Madbouly, M.D. (University of Alexandria, Alexandria, Egypt)
Koray Topgöl, M.D. (Anatolian Health Center, Kocaeli)
Levhi Akın, M.D. (Liv Hospital, İstanbul)
Liliana G. Bordeianou, M.D. (Massachusetts General Hospital, Harvard Medical School, Boston, USA)
Maria Cristina Sartor, M.D. (Pontificia Universidade Catolica do Parana, Parana, Brazil)
Mark Wong, M.D. (National University of Singapore, Singapore)
Massarat Zutshi, M.D. (Cleveland Clinic, Ohio, USA)
Mehmet Mihmanlı, M.D. (Etfal Training and Research Hospital, İstanbul)
Mehrdad Bohlooli, M.D. (Jam Hospital, Tehran, Iran)
Melih Paksoy, M.D. (İstanbul University Cerrahpaşa Faculty of Medicine, İstanbul)
Metin Ertem, M.D. (Acıbadem University Faculty of Medicine, İstanbul)
Mustafa Ateş, M.D. (İnönü University Faculty of Medicine, Malatya)
Mustafa Korkut, M.D. (Ege University Faculty of Medicine, İzmir)
Mustafa Öncel, M.D. (Medipol University Faculty of Medicine, İstanbul)
Neriman Şengül, M.D. (İzzet Baysal University Faculty of Medicine, Bolu)
Neşet Köksal, M.D. (Ümraniye Training and Research Hospital, İstanbul)
Nihat Yavuz, M.D. (Acıbadem University Faculty of Medicine, İstanbul)
Nuri Okkabaz, M.D. (Bağcılar Training and Research Hospital, İstanbul)
Oktar Asoğlu, M.D. (Bogazici Clinical Science Academy, İstanbul)
Ömer Alabaz, M.D. (Çukurova University Faculty of Medicine, Adana)
Ömer Topçu, M.D. (Cumhuriyet University Faculty of Medicine, Sivas)
Övünç Bardakçıoğlu, M.D. (University of Nevada School of Medicine, Nevada, USA)
Pars Tunçyürek, M.D. (Adnan Menderes University Faculty of Medicine, Aydın)
Paul Antoine Lehur, M.D. (University Hospital of Nantes, Nantes, France)
Robert D. Madoff, M.D. (University of Minnesota, Minneapolis, USA)
Sabri Ergüney, M.D. (İstanbul University Cerrahpaşa Faculty of Medicine, İstanbul)
Sadık Yıldırım, M.D. (Kolan Hastanesi, İstanbul)
Sedar Yüceyar, M.D. (İstanbul University Cerrahpaşa Faculty of Medicine, İstanbul)
Selçuk Atamanalp, M.D. (Atatürk University Faculty of Medicine, Erzurum)
Selman Sökmen, M.D. (Dokuz Eylül University Faculty of Medicine, İzmir)
Seyid Vahid Hosseini, M.D. (Shiraz University of Medical Sciences, Shiraz, Iran)
Sezai Demirbaş, M.D. (TOBB ETU Hospital, Ankara)
Soren Laurberg, M.D. (Aarhus University Hospital, Aarhus, Denmark)
Sümer Yamaner, M.D. (Florence Nightingale Hospitals, İstanbul)
Süphan Ertürk, M.D. (İstanbul University Cerrahpaşa Faculty of Medicine, İstanbul)
Tan Kok Yang, M.D. (National University of Singapore, Singapore)
Tarık Akçal, Florence Nightingale Hospital, İstanbul
Tayfun Karahasanoğlu, M.D. (Acıbadem University Faculty of Medicine, İstanbul)
Tuncay Yılmazlar, M.D. (Uludağ University Faculty of Medicine, Bursa)
Turgut İpek, M.D. (Medical Park Hospitals, İstanbul)
Türker Bulut, M.D. (İstanbul University Faculty of Medicine, İstanbul)
Uğur Sungurtekin, M.D. (Pamukkale University Faculty of Medicine, Denizli)
Yılmaz Büyükcü, M.D. (İstanbul University Faculty of Medicine, İstanbul)
Yunus Emre Altuntaş, M.D. (Lütfi Kırdar Training and Research Hospital, İstanbul)
Yüksel Altınel, M.D. (Harvard Medical School, Boston, USA)

Turkish Journal of COLORECTAL DISEASE



Aims and Scope

Turkish Journal of Colorectal Disease is an official journal of the Turkish Society of Colon and Rectal Surgery to provide epidemiologic, pathologic, diagnostic and therapeutic studies relevant to the management of small intestine, colon, rectum, anus and pelvic floor diseases. It was launched in 1991. Although there were temporary interruptions in the publication of the journal due to various challenges, the Turkish Journal of Colorectal Disease has been published continually from 2007 to the present. It is published quarterly (March, June, September and December) as hardcopy and an electronic journal at <http://www.turkishjcrd.com/>. The target audience of Turkish Journal of Colorectal Disease includes surgeons, pathologists, oncologists, gastroenterologists and health professionals caring for patients with a disease of the colon and rectum.

The Turkish name of the journal was formerly Kolon ve Rektum Hastalıkları Dergisi and the English name of the journal was formerly Journal of Diseases of the Colon and Rectum.

Turkish Journal of Colorectal Disease is indexed in TÜBİTAK/ULAKBİM, Directory of Open Access Journals (DOAJ), CINAHL Ultimate, British Library, Root Indexing, Academic Keys, Idealonline, Gale/Cengage Learning, Index Copernicus, Turkish Citation Index, Hinari, GOALI, ARDI, OARE and TurkMedline.

The aim of Turkish Journal of Colorectal Disease is to publish original research papers of the highest scientific and clinical value at an international level. Furthermore, review articles, case reports, technical notes, letters to the editor, editorial comments, educational contributions and congress/meeting announcements are released.

Turkish Journal of Colorectal Disease is an independent open access peer-reviewed international journal printed in Turkish and English languages. Manuscripts are reviewed in accordance with "double-blind peer review" process for both referees and authors. The Editorial Board of the Turkish Journal of Colorectal Disease endorses the editorial policy statements approved by the WAME Board of Directors. The journal is in compliance with the uniform requirements for manuscripts submitted to biomedical journals published by the International Committee of Medical Journal Editors (NEJM 1997;336:309-315, updated 2001).

Open Access Policy

This journal provides immediate open access to its content on the principle that making research freely available to the public supports a greater global exchange of knowledge. Open Access Policy is based on rules of Budapest Open Access Initiative (BOAI) <http://www.budapestopenaccessinitiative.org/>.

This journal is licensed under a Creative Commons 3.0 International License.

Permission Requests

Permission required for use any published under CC-BY-NC license with commercial purposes (selling, etc.) to protect copyright owner and author rights). Reproduction and reproduction of images or tables in any published material should be done with proper citation of source providing authors names; article title; journal title; year (volume) and page of publication; copyright year of the article.

Instructions for Authors

Instructions for authors are published in the journal and at www.turkishjcrd.com

Material Disclaimer

Authors are responsible for the manuscripts they publish in Turkish Journal of Colorectal Disease. The editor, editorial board, and publisher do not accept any responsibility for published manuscripts.

If you use a table or figure (or some data in a table or figure) from another source, cite the source directly in the figure or table legend.

The journal is printed on acid-free paper.

Financial expenses of the journal are covered by Turkish Society of Colon and Rectal Surgery.

Editorial Policy

Following receipt of each manuscript, a checklist is completed by the Editorial Assistant. The Editorial Assistant checks that each manuscript contains all required components and adheres to the author guidelines, after which time it will be forwarded to the Editor in Chief. Following the Editor in Chief's evaluation, each manuscript is forwarded to the Associate Editor, who in turn assigns reviewers. Generally, all manuscripts will be reviewed by at least three reviewers selected by the Associate Editor, based on their relevant expertise. Associate editor could be assigned as a reviewer along with the reviewers. After the reviewing process, all manuscripts are evaluated in the Editorial Board Meeting.

Turkish Journal of Colorectal Disease's editor and Editorial Board members are active researchers. It is possible that they would desire to submit their manuscript to the Turkish Journal of Colorectal Disease. This may be creating a conflict of interest. These manuscripts will not be evaluated by the submitting editor(s). The review process will be managed and decisions made by editor-in-chief who will act independently. In some situation, this process will be overseen by an outside independent expert in reviewing submissions from editors.

Subscription Information

Turkish Journal of Colorectal Disease is sent free - of - charge to members of Turkish Society of Colon and Rectal Surgery and libraries in Turkey and abroad. All published volumes are available in full text free-of-charge online at www.turkishjcrd.com

Address: Latilokum Sok. Alphan İşhanı No: 3 Kat: 2, Şişli, İstanbul, Türkiye

Telephone: +90 (212) 356 01 75-76-77

Gsm: +90 (532) 300 72 36

Fax: +90 (212) 356 01 78

Online Manuscript Submission: www.journalagent.com/krhd

Web page: www.turkishjcrd.com

E-mail: info@turkishjcrd.com

Advertisement / Publisher Corresponding Address

For requests concerning advertising, please contact the Publisher:

Galenos Yayınevi Tic. Ltd. Şti.

Address: Molla Garani Cad. 22/2 34093 Fındıkzade-İstanbul-Türkiye

Telephone: +90 (212) 621 99 25

Fax: +90 (212) 621 99 27

Web page: www.galenos.com.tr

E-mail: info@galenos.com.tr

Turkish Journal of COLORECTAL DISEASE



Amaç ve Kapsam

Türk Kolon ve Rektum Hastalıkları Dergisi, Türk Kolon ve Rektum Cerrahi Derneği'nin resmi dergisidir. Bu dernek; ince barsak, kolon, rektum, anüs ve pelvik taban hastalıkları gibi hastalıkların yönetimi ile ilişkili epidemiyolojik patolojik, tanısal ve tedavi edici çalışmalar yapar. Derneğimiz 1991'de kurulmuştur. Çeşitli zorluklar nedeniyle geçici aksaklıklar olsa da Türk Kolon ve Rektum Hastalıkları Dergisi 2007'den bu yana aralıksız olarak basılmaktadır ve 3 ayda bir olmak üzere (Mart, Haziran, Eylül, Aralık) basılı dergi ve elektronik olarak (<http://www.turkishjcrd.com/>) yayımlanır.

Derginin hedef kitesini; cerrahlar, patoloğlar, onkologlar, gastroenterologlar ve kolorektal hastalarına hizmet veren profesyoneller oluşturur. Derginin amacı; uluslararası düzeyde en yüksek bilimsel ve klinik değeri olan orijinal çalışmalarını yayımlamaktır. Bunlara ek olarak derleme (review) makaleleri, olgu sunumları, teknik notlar, editöre mektuplar, editöryal yorumlar, eğitim yazıları ve kongre/toplantı duyuruları yer almaktadır.

Derginin Türkçe eski adı; Kolon ve Rektum Hastalıkları Dergisi ve İngilizce eski adı; Journal of Diseases of the Colon and Rectum'dur.

Kolon ve Rektum Hastalıkları Dergisi, TÜBİTAK/ULAKBİM, Directory of Open Access Journals (DOAJ), CINAHL Ultimate, British Library, Root Indexing, Academic Keys, Ideonline, Gale/Cengage Learning, Index Copernicus, Türk Atf Dizini, Hinari, GOALI, ARDI, OARE ve TürkMedline'de indekslenmektedir..

Türk Kolon ve Rektum Hastalıkları Dergisi, İngilizce ve Türkçe olarak yayımlanan; bağımsız, hakemli, uluslararası bir dergidir. Eserler, hem hakemler hem de otörler tarafından "çift kör hakem denetimi (peer review)" yöntemi ile değerlendirilir. Türk Kolon ve Rektum Hastalıkları Dergisi'nin Editör Kurulu, World Association of Medical Editors (WAME) politikalarına bağlı olarak yürütülmektedir. Bu dergi, Uluslararası Tıp Dergisi Editörler Komitesi (NEJM 1997;336:309-315, updated 2001) tarafından bildirilen, biyomedikal dergilere gönderilen makalelerin uyması gereken standartlara uygunluk göstermektedir.

Açık Erişim Politikası

Bu dergi bilginin yer değiştirmesi ve topluluğu içinde bilgiye özgürce ulaşma olanağı sağlamak üzere açık erişime imkan vermektedir. Açık Erişim İlkesi "Budapeşte Açık Erişim Girişimi (BOAI)" <http://www.budapestopenaccessinitiative.org/> kurallarına dayanmaktadır.

Bu dergi Creative Commons 3.0 Uluslararası Lisansı ile lisanslanmıştır.

İzinler

Ticari amaçlarla CC-BY-NC lisansı altında yayımlanan her hangi bir kullanım (satış vb.) telif hakkı sahibi ve yazar haklarının korunması için izin gereklidir. Yayımlanan herhangi bir materyalde figure veya tabloların yeniden yayımlanması ve çoğaltılması, kaynağın başlık ve makalelerin yazarları ile doğru alıntılanmasıyla yapılmalıdır.

Derginin mali giderleri Türk Kolon ve Rektum Cerrahi Derneği tarafından karşılanmaktadır.

Yazarlar için Kılavuz

Yazarlar için kılavuz hem yayımlanan dergide hem de "<http://www.turkishjcrd.com/>" web sayfasında bulunmaktadır.

Telif Hakkı Devri

Yazarlar Türk Kolon ve Rektum Hastalıkları Dergisi'nde yayınladıkları yazılardan kendileri sorumludurlar. Editör, editör kurulu ve yayıncı hiçbir sorumluluk kabul etmemektedir. Başka bir kaynaktan tablo ya da figür (veya tablo/figürden bir veri) kullandıysanız, direkt olarak tablo ya da figürü kaynak gösteriniz.

Dergi asitsiz kağıda basılmaktadır.

Derginin mali giderleri Türk Kolon ve Rektum Cerrahi Derneği tarafından karşılanmaktadır.

Editöryal Politika

Her yazının alınmasını takiben, bir kontrol listesi Editör Yardımcısı tarafından tamamlanır.

Editör yardımcısı, her yazıyı gerekli öğeleri sağladığı ve yazar kılavuzuna uyumu açısından kontrol eder, ardından editöre iletir. Editör değerlendirmesinin ardından her bir yazı için editör yardımcısı tarafından gözlemciler (reviewers) belirlenir. Genelde, her bir yazıyı ilgili uzmanlıkları göz önüne alınarak atanmış en az 3 gözlemci inceler. Yardımcı editör de diğer gözlemcilerle birlikte gözlemci olarak atanabilir. Gözlemci incelemesinin ardından yazılar editör kurul toplantısında değerlendirilir.

Türk Kolon ve Rektum Hastalıkları Dergisi'nin editör ve editör kurulu üyeleri aktif araştırmacılar. Kendi araştırmalarının da Türk Kolon ve Rektum Hastalıkları Dergisi'nde yayınlanmasını pek ala arzu edebilirler. Bu durum çıkar sorunları doğurabilir. Bu yazılar, yazıyı yazan editör(ler) tarafından değerlendirilemez. Bu gibi durumlarda bu süreç, (editörlerin yazı başvurularında) yazıların uzman olan bağımsız kişiler tarafından incelenmesiyle aşılabılır.

Abonelik Bilgileri

Türk Kolon ve Rektum Hastalıkları Dergisi, Türk Kolon ve Rektum Cerrahisi Derneği üyelerine, Dünya'da ve Türkiye'deki kütüphanelere ücretsiz dağıtılmaktadır. Yayımlanmış tüm sayılar ücretsiz olarak şu linkte mevcuttur (<http://www.turkishjcrd.com/>).

Adres: Latilokum Sok. Alphan İşhanı No: 3 Kat: 2, Şişli, İstanbul, Türkiye

Telefon: +90 212 356 01 75-76- 77

GSM: +90 532 300 72 36

Faks: +90 212 356 01 78

Online Makale Gönderme: www.journalagent.com/krhd

Web sayfası: www.turkishjcrd.com

E-posta: info@turkishjcrd.com

Reklam-Duyuru / Yayınevi Yazışma Adresi

Talepleriniz için lütfen yayıncı ile iletişime geçiniz.

Galenos Yayınevi Tic. Ltd. Şti.

Molla Gürani Mah. Kaçamak Sk. No:21 34093 Fındıkzade-İstanbul-Türkiye

Telefon: +90 212 621 99 25 - Faks: +90 212 621 99 27

E-posta: info@galenos.com.tr

Web sayfası: www.galenos.com.tr



Turkish Journal of COLORECTAL DISEASE



Instruction for Authors

GENERAL INFORMATION

Turkish Journal of Colorectal Disease (TJCD) is the journal of Turkish Society of Colon and Rectal Surgery. The mission of the Journal is to advance knowledge of disorders of the small intestine, colon, rectum, anus and pelvic floor. It publishes invited review articles, research articles, brief reports and letters to the editor, and case reports that are relevant to the scope of the journal, on the condition that they have not been previously published elsewhere. Basic science manuscripts, such as randomized, cohort, cross-sectional, and case control studies, are given preference. Invited reviews will be considered for peer review from known experts in the area.

Manuscripts should be prepared according to ICMJE guidelines (www.icmje.org). All manuscripts are subject to editorial revision to ensure they conform to the style adopted by the journal. There is a double blind kind of reviewing system.

Reviewed and accepted manuscripts are translated from Turkish to English by the Journal through a professional translation service. Prior to printing, the translations are submitted to the authors for approval or correction requests, to be returned within 7 days. If no response is received from the corresponding author within this period, the translation is checked and approved by the editorial board.

Accepted manuscripts are published in both Turkish and English languages.

All manuscripts submitted to the Turkish Journal of Colorectal Disease are screened for plagiarism using the 'iThenticate' software. Results indicating plagiarism may result in manuscripts being returned or rejected.

Turkish Journal of Colorectal Disease does not charge any article submission or processing charges.

The abbreviation of the Turkish Journal of Colorectal Disease is "TJCD", however, it should be denoted as "Turk J Colorectal Dis" when referenced.

EDITORIAL POLICIES

All manuscripts will be evaluated by the scientific board for their scientific contribution, originality and content. Authors are responsible for the accuracy of the data. The journal retains the right to make appropriate changes on the grammar and language of the manuscript. When suitable the manuscript will be sent to the corresponding author for revision. The manuscript, when published, will become the property of the journal and copyright will be taken out in the name of the journal

"Turkish Journal of Colorectal Disease". Articles previously published in any language will not be considered for publication in the journal. Authors cannot submit the manuscript for publication in another journal. All changes in the manuscript will be made after obtaining written permission of the author and the publisher. Full text of all articles can be downloaded at the web site of the journal www.journalagent.com/krhd.

AUTHOR GUIDELINES

Forms Required with Submission:

Copyright Transfer Statement
Disclosure Statement
Cover Letter

Manuscript Submission Guidelines

Manuscript Preparation Guidelines
Text Formatting
Title Page
Article Types
Original Articles
Invited Review Articles
Case Reports
Technical Notes
Letters to Editor
Editorial Comments
Ethical Responsibilities of Authors
Research Involving Human Participants and/or Animals
Informed Consent
Payment

Forms Required with Submission

Copyright Transfer Statement

The scientific and ethical liability of the manuscripts belongs to the authors and the copyright of the manuscripts belongs to the Turkish Journal of Colorectal Disease. Authors are responsible for the contents of the manuscript and accuracy of the references. All manuscripts submitted for publication must be accompanied by the Copyright Transfer Form [copyright transfer]. Once this form, signed by all the authors, has been submitted, it is understood that neither the manuscript nor the data it contains have been submitted elsewhere or previously published and authors declare the statement of scientific contributions and responsibilities of all authors.

Disclosure Statement

Conflicts of interest: Authors must state all possible conflicts of interest in the manuscript, including financial, consultant, institutional and other relationships that might lead to bias or a conflict of interest. If there is no conflict of interest, this should also be explicitly stated as none declared. All sources of funding should be acknowledged in the manuscript. All relevant conflicts of interest and sources of funding should be included on the title page of the manuscript with the heading "Conflicts of Interest and Source of Funding:"

Cover Letter

In the cover letter the authors should state if any of the material in the manuscript is submitted or planned for publication elsewhere in any form including electronic media. A written statement indicating whether or not "Institutional Review Board" (IRB) approval was obtained or equivalent guidelines followed in accordance with the Helsinki Declaration of

2013 update on human experimentation must be stated; if not, an explanation must be provided. The cover letter must contain address, telephone, fax and the e-mail address of the corresponding author.

Manuscript Submission Guidelines

All manuscripts should be submitted via the online submission system. Authors are encouraged to submit their manuscripts via the internet after logging on to the web site www.journalagent.com/krhd.

The ORCID (Open Researcher and Contributor ID) number of the correspondence author should be provided while sending the manuscript. A free registration can create at <http://orcid.org>.

Online Submission

Only online submissions are accepted for rapid peer-review and to prevent delay in publication. Manuscripts should be prepared as word document (*.doc) or rich text format (*.rtf). After logging on to the web www.journalagent.com/krhd double click the "submit an article" icon. All corresponding authors should be provided a password and an username after providing the information needed. After logging on the article submission system with your own password and username please read carefully the directions of the system to provide all needed information in order not to delay the processing of the manuscript. Attach the manuscript, all figures, tables and additional documents. Please also attach the cover letter with "Assignment of Copyright and Financial Disclosure" forms.

Manuscript Preparation Guidelines

Turkish Journal of Colorectal Disease follows the "Uniform Requirements for Manuscripts Submitted to Biomedical Journals" (International Committee of Medical Journal Editors: Br Med J 1988;296:401-5).

Upon submission of the manuscript, authors are to indicate the type of trial/research and statistical applications following "Guidelines for statistical reporting in articles for medical journals: amplifications and explanations" (Bailar JC III, Mosteller F. Ann Intern Med 1988;108:266-73).

Preparation of research articles, systematic reviews and meta-analyses must comply with study design guidelines:

CONSORT statement for randomized controlled trials (Moher D, Schultz KF, Altman D, for the CONSORT Group. The CONSORT statement revised recommendations for improving the quality of reports of parallel group randomized trials. JAMA 2001; 285:1987-91) (<http://www.consort-statement.org/>);

PRISMA statement of preferred reporting items for systematic reviews and meta-analyses (Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group. Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 2009; 6(7): e1000097.) (<http://www.prisma-statement.org/>);

STARD checklist for the reporting of studies of diagnostic accuracy (Bossuyt PM, Reitsma JB, Bruns DE, Gatsonis CA,

Turkish Journal of COLORECTAL DISEASE



Instruction for Authors

Glasziou PP, Irwig LM, et al., for the STARD Group. Towards complete and accurate reporting of studies of diagnostic accuracy: the STARD initiative. *Ann Intern Med* 2003;138:40-4. (<http://www.stard-statement.org/>);

STROBE statement, a checklist of items that should be included in reports of observational studies (<http://www.strobe-statement.org/>);

MOOSE guidelines for meta-analysis and systemic reviews of observational studies (Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting Meta-analysis of observational Studies in Epidemiology (MOOSE) group. *JAMA* 2000; 283: 2008-12).

Text Formatting

Manuscripts should be submitted in Word.

Use a normal, plain font (e.g., 10-point Times Roman) for text.

Use the automatic page numbering function to number the pages.

Do not use field functions.

Use tab stops or other commands for indents, not the space bar.

Use the table function, not spreadsheets, to make tables.

Save your file in docx format (Word 2007 or higher) or doc format (older Word versions).

Title Page

All manuscripts, regardless of article type, should start with a title page, containing:

The title of the article;

The short title of the article

The initials, names and qualifications of each author;

The main appointment of each author;

The name(s) of the institution(s) of each author;

The name and email address of the corresponding author;

Full disclosures of potential conflicts of interest on the part of any named author, or a statement confirming that there are no conflicts of interest;

The word count excluding abstract, references, tables, figures and legends;

The place and date of scientific meeting in which the manuscript was presented and it's abstract published in the abstract book, if applicable.

Article Types

Original Articles

This category includes original research including both clinical and basic science submissions. The work must be original and neither published, accepted, or submitted for publication elsewhere. Any related work, either SUBMITTED, in press, or published from any of the authors should be clearly cited and referenced.

All clinical trials must be registered in a public trials registry that is acceptable to the International Committee of Medical

Journals Editors (ICMJE). Go to (<http://www.icmje.org/faq.html>). Authors of randomized controlled trials must adhere to the CONSORT guidelines, available at: www.consort-statement.org, and provide both a CONSORT checklist and flow diagram. We require that you choose the MS Word template at www.consort-statement.org for the flow chart and cite/upload it in the manuscript as a figure. In addition, submitted manuscripts must include the unique registration number in the Abstract as evidence of registration.

All authors are expected to abide by accepted ethical standards for human and animal investigation. In studies that involve human subjects or laboratory animals, authors must provide an explicit statement in Materials and Methods that the experimental protocol was approved by the appropriate institutional review committee and meets the guidelines of their responsible governmental agency. In the case of human subjects, informed consent, in addition to institutional review board approval, is required.

Original Articles should not exceed 3000 words (excluding abstract, references, tables, figures and legends) and four illustrations.

Original Articles should be organized as follows:

Abstract: The abstract must contain fewer than 250 words and should be structured as follows:

Aim: What was the purpose of the study?

Method: A brief description of the materials - patients or subjects (i.e. healthy volunteers) or materials (animals) - and methods used.

Results: What were the main findings?

Conclusion: What are the main conclusions or implications of the study?

Keywords: Below the abstract provide up to 6 key words or short phrases. Do not use abbreviations as keywords.

Introduction: State concisely the purpose and rationale for the study and cite only the most pertinent references as background.

Materials and Methods: Describe your selection of the observational or experimental subjects clearly (patients or experimental animals, including controls). Provide an explicit statement that the experimental protocols were approved by the appropriate institutional review committee and meet the guidelines of the responsible governmental agency. In the case of human subjects, state explicitly those subjects have provided informed consent. Identify the methods, apparatus/product** (with manufacturer's name and address in parentheses), and procedures in sufficient detail to allow other workers to reproduce the results. Give references to established methods, including statistical methods; provide references and brief descriptions of methods that have been published but are not well known, describe substantially modified methods, including statistical methods, give reasons for using them, and evaluate their limitations;

Results: Present the detailed findings supported with statistical methods. Figures and tables should supplement, not duplicate the text; presentation of data in either one or the other will suffice. Emphasize only your important observations; do not compare your observations with those of others. Such comparisons and comments are reserved for the discussion section.

Discussion: State the importance and significance of your findings but do not repeat the details given in the Results section. Limit your opinions to those strictly indicated by the facts in your report. Compare your finding with those of others. No new data are to be presented in this section.

Acknowledgments: Only acknowledge persons who have made substantive contributions to the study. Authors are responsible for obtaining written permission from everyone acknowledged by name because readers may infer their endorsement of the data and conclusions. Begin your text of the acknowledgment with, "The authors thank...".

Authorship Contributions: The journal follows the recommendations of the ICMJE for manuscripts submitted to biomedical journals. According to these, authorship should be based on the following four criteria:

Substantial contributions to the conception or design of the work; or the acquisition, analysis, or interpretation of data for the work; and

Drafting the work or revising it critically for important intellectual content; and

Final approval of the version to be published; and

Agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

All other contributors to the paper should be credited in the 'Acknowledgments' section.

References: The author should number the references in Arabic numerals according to the citation order in the text. Put reference numbers in parenthesis in superscript at the end of citation content or after the cited author's name. Use the form of "Uniform Requirements for manuscript abbreviations in Turk Bilim Terimleri" (<http://www.bilimterimleri.com>).

Journal titles should conform to the abbreviations used in "Cumulated Index Medicus".

Journals; Last name(s) of the author(s) and initials, article title, publication title and its original abbreviation, publication date, volume, the inclusive page numbers.

Example: 1. Dilaveris P, Batchvarov V, Gialafos J, Malik M. Comparison of different methods for manual P wave duration measurement in 12-lead electrocardiograms. *Pacing Clin Electrophysiol* 1999;22:1532-1538.

Book chapter; Last name(s) of the author(s) and initials, chapter title, book editors, book title, edition, place of publication, date of publication and inclusive page numbers of the extract cited.

Turkish Journal of COLORECTAL DISEASE



Instruction for Authors

Example: 1. Schwartz PJ, Priori SG, Napolitano C. The Long QT Syndrome. In: Zipes DP, Jalife J, eds. Cardiac Electrophysiology. From Cell to Bedside. Philadelphia; WB Saunders Co. 2000:597-615.

Tables: All tables are to be numbered using Arabic numerals. Tables should always be cited in text in consecutive numerical order. For each table, please supply a table caption (title) explaining the components of the table. Identify any previously published material by giving the original source in the form of a reference at the end of the table caption. Footnotes to tables should be indicated by superscript lower-case letters (or asterisks for significance values and other statistical data) and included beneath the table body.

Figures: Figures should work under "Windows". Color figures or grayscale images must be at least 300 dpi. Figures using "*.tiff", "*.jpg" or "*.pdf" should be saved separate from the text. All figures should be prepared on separate pages. They should be numbered in Arabic numerals. Each figure must have an accompanying legend defining abbreviations or symbols found in the figure. Figures could be submitted at no additional cost to the author.

Units of Measurement and Abbreviations: Units of measurement should be in Système International (SI) units. Abbreviations should be avoided in the title. Use only standard abbreviations. If abbreviations are used in the text, they should be defined in the text when first used.

Permissions: Authors wishing to include figures, tables, or text passages that have already been published elsewhere are required to obtain permission from the copyright owner(s) and to include evidence that such permission has been granted when submitting their papers. Any material received without such evidence will be assumed to originate from the authors.

Invited Review Articles

Abstract length: Not to exceed 250 words.

Article length: Not to exceed 4000 words.

Reference Number: Not to exceed 100 references.

Reviews should include a conclusion, in which a new hypothesis or study about the subject may be posited. Do not publish methods for literature search or level of evidence. Authors who will prepare review articles should already have published research articles on the relevant subject. The study's new and important findings should be highlighted and interpreted in the Conclusion section. There should be a maximum of two authors for review articles.

Case Reports

Abstract length: Not to exceed 100 words.

Article length: Not to exceed 1000 words.

Reference Number: Not to exceed 15 references.

Case Reports should be structured as follows:

Abstract: An unstructured abstract that summarizes the case.

Introduction: A brief introduction (recommended length: 1-2 paragraphs).

Case Report: This section describes the case in detail, including the initial diagnosis and outcome.

Discussion: This section should include a brief review of the relevant literature and how the presented case furthers our understanding to the disease process.

References: See under 'References' above.

Acknowledgments.

Tables and figures.

Technical Notes

Abstract length: Not to exceed 250 words.

Article length: Not to exceed 1200 words.

Reference Number: Not to exceed 15 references.

Technical Notes include description of a new surgical technique and its application on a small number of cases. In case of a technique representing a major breakthrough one case will suffice. Follow-up and outcome need to be clearly stated.

Technical Notes should be organized as follows:

Abstract: Structured "as above mentioned".

Indications

Method

Comparison with other methods: advantages and disadvantages, difficulties and complications.

References, in Vancouver style (see under 'References' above).

Acknowledgments.

Tables and figures: Including legends.

Letters to the Editor

Article length: Not to exceed 500 words.

Reference Number: Not to exceed 10 references

We welcome correspondence and comment on articles published in Turkish Journal of Colorectal Disease. No abstract is required, but please include a brief title. Letters can include 1 figure or table.

Video Article

Article length: Not to exceed 500 words.

Reference Number: Not to exceed 5 references

Briefly summarize the case describing diagnosis, applied surgery technique and outcome. Represent all important aspects, i.e. novel surgery technique, with properly labelled and referred video materials. A standalone video vignette, describing a surgical technique or interesting case encountered by the authors.

Requirements: The data must be uploaded during submission with other files. The video should be no longer than 10 minutes in duration with a maximum file size of 350Mb and 'MOV, MPEG4, AVI, WMV, MPEGPS, FLV, 3GPP, WebM' format should be used. Documents that do not exceed 100 MB can be uploaded within the system. For larger video documents, please contact iletisim@galenos.com.tr All videos must include a narration in English. Reference must be used as it would be for a Figure or a Table. Example: ".....To accomplish this, we developed

a novel surgical technique (Video 1)." All names and institutions should be removed from all video materials. Video materials of accepted manuscripts will be published online.

Letters to the Editor

Article length: Not to exceed 500 words.

Reference Number: Not to exceed 10 references

We welcome correspondence and comment on articles published in Turkish Journal of Colorectal Disease. No abstract is required, but please include a brief title. Letters can include 1 figure or table.

Editorial Comments

Article length: Not to exceed 1000 words.

Reference Number: Not to exceed 10 references.

Editorials are exclusively solicited by the Editor. Editorials should express opinions and/or provide comments on papers published elsewhere in the same issue. A single author is preferred. No abstract is required, but please include a brief title. Editorial submissions are subject to review/request for revision, and editors retain the right to alter text style.

Ethics

This journal is committed to upholding the integrity of the scientific record. As a member of the Committee on Publication Ethics (COPE) the journal will follow the COPE guidelines on how to deal with potential acts of misconduct.

Authors should refrain from misrepresenting research results which could damage the trust in the journal, the professionalism of scientific authorship, and ultimately the entire scientific endeavor. Maintaining integrity of the research and its presentation can be achieved by following the rules of good scientific practice, which include:

The manuscript has not been submitted to more than one journal for simultaneous consideration.

The manuscript has not been published previously (partly or in full), unless the new work concerns an expansion of previous work (please provide transparency on the re-use of material to avoid the hint of text-recycling ("self-plagiarism").

A single study is not split up into several parts to increase the quantity of submissions and submitted to various journals or to one journal over time (e.g. "salami-publishing").

No data have been fabricated or manipulated (including images) to support your conclusions.

No data, text, or theories by others are presented as if they were the author's own ("plagiarism"). Proper acknowledgments to other works must be given (this includes material that is closely copied (near verbatim), summarized and/or paraphrased), quotation marks are used for verbatim copying of material, and permissions are secured for material that is copyrighted.

Important note: Turkish Journal of Colorectal Disease uses software (iThenticate) to screen for plagiarism.

Turkish Journal of COLORECTAL DISEASE



Instruction for Authors

Consent to submit has been received explicitly from all co-authors, as well as from the responsible authorities - tacitly or explicitly - at the institute/organization where the work has been carried out, before the work is submitted.

Authors whose names appear on the submission have contributed sufficiently to the scientific work and therefore share collective responsibility and accountability for the results.

In addition: Changes of authorship or in the order of authors are not accepted after acceptance of a manuscript.

Requesting to add or delete authors at revision stage, proof stage, or after publication is a serious matter and may be considered when justifiably warranted. Justification for changes in authorship must be compelling and may be considered only after receipt of written approval from all authors and a convincing, detailed explanation about the role/deletion of the new/deleted author. In case of changes at revision stage, a letter must accompany the revised manuscript. In case of changes after acceptance or publication, the request and documentation must be sent via the Publisher to the Editor-in-Chief. In all cases, further documentation may be required to support your request. The decision on accepting the change rests with the Editor-in-Chief of the journal and may be turned down. Therefore authors are strongly advised to ensure the correct author group, corresponding author, and order of authors at submission.

Upon request authors should be prepared to send relevant documentation or data in order to verify the validity of the results. This could be in the form of raw data, samples, records, etc.

If there is a suspicion of misconduct, the journal will carry out an investigation following the COPE guidelines. If, after investigation, the allegation seems to raise valid concerns, the accused author will be contacted and given an opportunity to address the issue. If misconduct has been established beyond reasonable doubt, this may result in the Editor-in-Chief's implementation of the following measures, including, but not limited to:

If the article is still under consideration, it may be rejected and returned to the author.

If the article has already been published online, depending on the nature and severity of the infraction, either an erratum will be placed with the article or in severe cases complete retraction of the article will occur. The reason must be given in the published erratum or retraction note.

The author's institution may be informed.

Editorial Comments

Article length: Not to exceed 1000 words.

Reference Number: Not to exceed 10 references.

Editorials are exclusively solicited by the Editor. Editorials should express opinions and/or provide comments on papers published elsewhere in the same issue. A single author is preferred. No abstract is required, but please include a brief title. Editorial submissions are subject to

review/request for revision, and editors retain the right to alter text style.

Ethics

This journal is committed to upholding the integrity of the scientific record. As a member of the Committee on Publication Ethics (COPE) the journal will follow the COPE guidelines on how to deal with potential acts of misconduct.

Authors should refrain from misrepresenting research results which could damage the trust in the journal, the professionalism of scientific authorship, and ultimately the entire scientific endeavor. Maintaining integrity of the research and its presentation can be achieved by following the rules of good scientific practice, which include:

The manuscript has not been submitted to more than one journal for simultaneous consideration.

The manuscript has not been published previously (partly or in full), unless the new work concerns an expansion of previous work (please provide transparency on the re-use of material to avoid the hint of text-recycling ("self-plagiarism").

A single study is not split up into several parts to increase the quantity of submissions and submitted to various journals or to one journal over time (e.g. "salami-publishing").

No data have been fabricated or manipulated (including images) to support your conclusions.

No data, text, or theories by others are presented as if they were the author's own ("plagiarism"). Proper acknowledgments to other works must be given (this includes material that is closely copied (near verbatim), summarized and/or paraphrased), quotation marks are used for verbatim copying of material, and permissions are secured for material that is copyrighted.

Important note: Turkish Journal of Colorectal Disease uses software (iThenticate) to screen for plagiarism.

Consent to submit has been received explicitly from all co-authors, as well as from the responsible authorities - tacitly or explicitly - at the institute/organization where the work has been carried out, before the work is submitted.

Authors whose names appear on the submission have contributed sufficiently to the scientific work and therefore share collective responsibility and accountability for the results.

In addition: Changes of authorship or in the order of authors are not accepted after acceptance of a manuscript.

Requesting to add or delete authors at revision stage, proof stage, or after publication is a serious matter and may be considered when justifiably warranted. Justification for changes in authorship must be compelling and may be considered only after receipt of written approval from all authors and a convincing, detailed explanation about the role/deletion of the new/deleted author. In case of changes at revision stage, a letter must accompany the revised manuscript. In case of changes after acceptance or publication, the request and documentation must be sent

via the Publisher to the Editor-in-Chief. In all cases, further documentation may be required to support your request. The decision on accepting the change rests with the Editor-in-Chief of the journal and may be turned down. Therefore authors are strongly advised to ensure the correct author group, corresponding author, and order of authors at submission.

Upon request authors should be prepared to send relevant documentation or data in order to verify the validity of the results. This could be in the form of raw data, samples, records, etc.

If there is a suspicion of misconduct, the journal will carry out an investigation following the COPE guidelines. If, after investigation, the allegation seems to raise valid concerns, the accused author will be contacted and given an opportunity to address the issue. If misconduct has been established beyond reasonable doubt, this may result in the Editor-in-Chief's implementation of the following measures, including, but not limited to:

If the article is still under consideration, it may be rejected and returned to the author.

If the article has already been published online, depending on the nature and severity of the infraction, either an erratum will be placed with the article or in severe cases complete retraction of the article will occur. The reason must be given in the published erratum or retraction note.

The author's institution may be informed.

Research Involving Human Participants and/or Animals

Statement of human rights: When reporting studies that involve human participants, authors should include a statement that the studies have been approved by the appropriate institutional and/or national research ethics committee and have been performed in accordance with the ethical standards as laid down in the 1964 Declaration of Helsinki and its later amendments or comparable ethical standards.

If doubt exists whether the research was conducted in accordance with the 1964 Helsinki Declaration or comparable standards, the authors must explain the reasons for their approach, and demonstrate that the independent ethics committee or institutional review board explicitly approved the doubtful aspects of the study.

The following statements should be included in the text before the References section: Ethical approval: "All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards."

For retrospective studies, please add the following sentence: "For this type of study formal consent is not required."

Statement on the welfare of animals: The welfare of animals used for research must be respected. In experimental animal

Turkish Journal of COLORECTAL DISEASE



Instruction for Authors

studies, the authors should indicate that the procedures followed were in accordance with animal rights as per the Guide for the Care and Use of Laboratory Animals <http://oacu.od.nih.gov/regs/guide/guide.pdf> and they should obtain animal ethics committee approval. When reporting experiments on animals, authors should indicate whether the international, national, and/or institutional guidelines for the care and use of animals have been followed, and that the studies have been approved by a research ethics committee at the institution or practice at which the studies were conducted (where such a committee exists).

For studies with animals, the following statement should be included in the text before the References section:

Ethical approval: "All applicable international, national, and/or institutional guidelines for the care and use of animals were followed."

If applicable (where such a committee exists): "All procedures performed in studies involving animals were in accordance with the ethical standards of the institution or practice at which the studies were conducted."

If articles do not contain studies with human participants or animals by any of the authors, please select one of the following statements:

"This article does not contain any studies with human participants performed by any of the authors."

"This article does not contain any studies with animals performed by any of the authors."

"This article does not contain any studies with human participants or animals performed by any of the authors."

Informed Consent

All individuals have individual rights that are not to be infringed. Individual participants in studies have, for example, the right to decide what happens to the (identifiable) personal data gathered, to what they have said during a study or an interview, as well as to any photograph that was taken. Hence it is important that all participants gave their informed consent in writing prior to inclusion in the study. Identifying details (names, dates of birth, identity numbers and other information) of the participants that were studied should not be published in written descriptions, photographs, and genetic profiles unless the information is essential for scientific purposes and the participant (or parent or guardian if the participant is incapable) gave written informed consent for publication. Complete anonymity is difficult to achieve in some cases, and informed consent should be obtained if there is any doubt. For example, masking the eye region in photographs of participants is inadequate protection of anonymity. If identifying characteristics are altered to protect anonymity, such as in genetic profiles, authors should provide assurance that alterations do not distort scientific meaning.

The following statement should be included: Informed Consent: "Informed consent was obtained from all individual participants included in the study."

If identifying information about participants is available in the article, the following statement should be included:

"Additional informed consent was obtained from all individual participants for whom identifying information is included in this article."

Payment

Turkish Journal of Colorectal Disease does not charge any article submission or processing charges.

THE REVIEW PROCESS

Each manuscript submitted to The Turkish Journal of Colorectal Disease is subject to an initial review by the editorial office in order to determine if it is aligned with the journal's aims and scope, and complies with essential requirements. Manuscripts sent for peer review will be assigned to one of the journal's associate editors that has expertise relevant to the manuscript's content. All accepted manuscripts are sent to a statistical and English language editor before publishing. Once papers have been reviewed, the reviewers' comments are sent to the Editor, who will then make a preliminary decision on the paper. At this stage, based on the feedback from reviewers, manuscripts can be accepted, rejected, or revisions can be recommended. Following initial peer-review, articles judged worthy of further consideration often require revision. Revised manuscripts generally must be received within 2 months of the date of the initial decision. Extensions must be requested from the Associate Editor at least 2 weeks before the 2-month revision deadline expires; The Turkish Journal of Colorectal Disease will reject manuscripts that are not received within the 3-month revision deadline. Manuscripts with extensive revision recommendations will be sent for further review (usually by the same reviewers) upon their re-submission. When a manuscript is finally accepted for publication, the Technical Editor undertakes a final edit and a marked-up copy will be e-mailed to the corresponding author for review and to make any final adjustments.

REVISIONS

When submitting a revised version of a paper, the author must submit a detailed "Response to the reviewers" that states point by point how each issue raised by the reviewers has been covered and where it can be found (each reviewer's comment, followed by the author's reply and line numbers where the changes have been made) as well as an annotated copy of the main document. Revised manuscripts must be submitted within 30 days from the date of the decision letter. If the revised version of the manuscript is not submitted within the allocated time, the revision option may be canceled. If the submitting author(s) believe that additional time is required, they should request this extension before the initial 30-day period is over.

ENGLISH LANGUAGE EDITING

All manuscripts are professionally edited by an English language editor prior to publication.

AFTER ACCEPTANCE

All accepted articles are technically edited by one of the Editors. On completion of the technical editing, the article will be sent to the production department and published online as a fully citable Accepted Article within about one week.

Copyright Transfer

Authors will be asked to transfer copyright of the article to the Publisher (or grant the Publisher exclusive publication and dissemination rights). This will ensure the widest possible protection and dissemination of information under copyright laws.

Color Illustrations

Publication of color illustrations is free of charge.

Proof Reading

The purpose of the proof is to check for typesetting or conversion errors and the completeness and accuracy of the text, tables and figures. Substantial changes in content, e.g., new results, corrected values, title and authorship, are not allowed without the approval of the Editor.

After online publication, further changes can only be made in the form of an Erratum, which will be hyperlinked to the article.

ONLINE EARLY

The Turkish Journal of Colorectal Disease publishes abstracts of accepted manuscripts online in advance of their publication in print. Once an accepted manuscript has been edited, the authors have submitted any final corrections, and all changes have been incorporated, the manuscript will be published online. At that time the manuscript will receive a Digital Object Identifier (DOI) number. Both forms can be found at www.journalagent.com/krhd. Authors of accepted manuscripts will receive electronic page proofs directly from the printer, and are responsible for proofreading and checking the entire manuscript, including tables, figures, and references. Page proofs must be returned within 48 hours to avoid delays in publication.

CORRESPONDENCE

All correspondences can be done to the following postal address or to the following e-mail address, where the journal editorial resides:

Address: Latilokum Sok. Alphan İşhanı No:3 Kat:2 Mecidiyeköy-Şişli-İstanbul- Turkey

Phone: +90 (212) 356 01 75-76-77

Gsm: +90 (532) 300 72 36

Fax: +90 (212) 356 01 78

Online Manuscript: www.journalagent.com/krhd

Web page: www.turkishjcrd.com

E-mail: info@turkishjcrd.com

Turkish Journal of COLORECTAL DISEASE



Yazarlara Bilgi

GENEL BİLGİ

Türk Kolon ve Rektum Hastalıkları Dergisi, Türk Kolon ve Rektum Cerrahisi Derneği'nin dergisidir. Derginin misyonu; ince bağırsak, kolon, rektum, anüs ve pelvik taban bozuklukları hakkındaki bilgiye katkı sağlamaktır. Dergi daha önce başka bir yerde yayınlanmamış olması koşuluyla, derginin kapsamı ile ilgili ve talep üzerine yazılan derleme makaleleri, araştırma makaleleri, kısa raporlar ve editöre mektuplar ve olgu sunumlarını yayınlamaktadır. Randomize, kohort, kesitsel ve vaka kontrol çalışmaları gibi temel bilim yazılarına öncelik verilir. Alanında bilinen uzmanlarca talep üzerine yazılan derlemeler dikkate alınacaktır.

Yazılar ICMJE yönergelerine göre (<http://www.icmje.org/>) hazırlanmalıdır. Tüm yazılar dergi tarafından benimsenen stile uygunluk sağlamak için editöryal kontrol ve düzeltmelere tabi tutulmaktadır. Derginin çift kör bir değerlendirme sistemi vardır. Değerlendirilen ve kabul edilen yayınlar Türkçeden İngilizceye veya İngilizceden Türkçeye derginin profesyonel çeviri hizmeti aracılığıyla tercüme edilir. Yayınlanmadan önce, çeviriler onay veya düzeltme istekleri için yazarlara gönderilir ve 7 gün içinde geri dönüş talep edilir. Bu süre içinde yanıt alınmazsa, çeviri kontrol ve yayın kurulu tarafından onaylanır.

Kabul edilen yayınlar hem Türkçe hem de İngilizce olarak yayınlanır.

Türk Kolon ve Rektum Hastalıkları Dergisi'ne gönderilen tüm yayınlar 'iThenticate' yazılımı kullanılarak intihal açısından taranır. İntihal saptanan durumlarda yayın iade veya reddedilir.

Türk Kolon ve Rektum Hastalıkları Dergisi, makale gönderme veya işlem ücreti adı altında herhangi bir ücret talep etmemektedir.

Türk Kolon ve Rektum Hastalıkları Dergisi'nin kısaltması "TJCD"dir, ancak, refere edildiğinde "Turk J Colorectal Dis" olarak kullanılmalıdır.

YAYIN POLİTİKASI

Tüm makaleler bilimsel katkıları, özgünlük ve içerikleri açısından bilimsel komite tarafından değerlendirilecektir. Yazarlar verilerinin doğruluğundan sorumludurlar. Dergi gerekli gördüğü yerlerde dil ve uygun değişiklik yapma hakkını saklı tutar. Gereğinde makale revizyon için yazara gönderilir. Dergide basılan yayının mali haline gelir ve telif hakkı "Türk Kolon ve Rektum Hastalıkları Dergisi" adına alınmış olur. Daha önce herhangi bir dilde yayınlanmış makaleler dergide yayınlanmak üzere kabul edilmeyecektir. Yazarlar bir başka dergide yayınlanmak üzere olan makaleyi teslim edemez. Tüm değişiklikler, yazar ve yayıncının yazılı izni alındıktan sonra yapılacaktır. Tüm makalelerin tam metinleri derginin www.journalagent.com/krhd web sitesinden indirilebilir.

YAZAR KILAVUZU

Makale gönderilirken sunulması gereken formlar:

Telif hakkı devir bildirimini

Açıklama bildirimini

Üst yazı

Makale Gönderme Kuralları

Makale Hazırlama Kuralları

Metin biçimlendirme

Giriş sayfası

Yayın tipleri

Orijinal Makaleler

Talepli derlemeler

Olgu sunumları

Teknik notlar

Editöre mektuplar

Editöryal Yorumlar

Yazarların Etik Sorumlulukları

İnsan katılımcılı araştırma ve/veya hayvan deneyleri

Bilgilendirilmiş Onam

Makale Gönderilirken Sunulması Gereken Formlar:

Telif Hakkı Devir Bildirimi

Yayınlann bilimsel ve etik sorumluluğu yazarlarına aittir. Yazıların telif hakkı ise Türk Kolon ve Rektum Hastalıkları Dergisi'ne aittir. Yazarlar yayınlann doğruluk ve içeriğinden ve kaynakların doğruluğundan sorumludur. Yayınlanmak üzere gönderilen tüm yayınlara Telif Hakkı Devir Formu (telif hakkı transferi) eşlik etmelidir. Tüm yazarlar tarafından imzalanarak gönderilen bu form ile yazarlar, ilgili yayının ve içerdiği datanın başka bir yayın organına gönderilmediğini veya başka bir dergide yayınlanmadığını beyan ederler. Ayrıca bu belge yazarların bilimsel katkı ve tüm sorumluluklarının ifadesidir.

Açıklama Bildirimi

Çıkar çatışmaları: Yazarlar, finansal, kurumsal, danışmanlık şeklinde ya da herhangi bir çıkar çatışmasına yol açabilecek başka ilişkiler de dahil olmak üzere yayındaki ilgili tüm olası çıkar çatışmalarını belirtmelidir. Herhangi bir çıkar çatışması yoksa da bu da açıkça belirtilmelidir. Tüm finansman kaynakları yazının içinde belirtilmelidir. Finansman kaynakları ve ilgili tüm çıkar çatışmaları yazının başlık sayfasında "Finansman ve Kaynak Çatışmaları:" başlığı ile yer almalıdır.

Üst Yazı

Yazarlar, yazının içinde malzemenin elektronik ortam da dahil olmak üzere herhangi bir başka bir yerde yayımlanmak üzere gönderilmediğini veya planlanmadığını üst yazıda belirtmelidir. Yine "Kurumsal Değerlendirme Kurulu" (KDK) onayı alınıp alınmadığı ve 2013 yılı Helsinki Bildirgesi'ne eşdeğer kılavuzların izlenip izlenmediği belirtilmelidir. Aksi takdirde, bir açıklama temin edilmelidir. Üst yazı; adres, telefon, faks ve ilgili yazının e-posta adresini içermelidir.

Makale Yazım Kuralları

Tüm makaleler online başvuru sistemi üzerinden teslim edilmelidir. Yazarlar web sitesi www.journalagent.com/krhd adresinde oturum açtıktan sonra internet üzerinden yazıların sunulmalıdır.

Makale gönderimi yapılırken sorumlu yazarın ORCID (Open Researcher ve Contributor ID) numarası belirtilmelidir. <http://orcid.org> adresinden ücretsiz olarak kayıt oluşturulabilir.

Online Başvuru

Gecikmeyi önlemek ve hızlı hakemlik için sadece çevrim içi gönderimler kabul edilir. Yazılar word belgesi (*.doc) veya zengin metin biçimi (*.rtf) olarak hazırlanmalıdır. www.journalagent.com/krhd adresinde web oturumu açtıktan sonra "Makale gönder" ikonuna tıklayın. Tüm yazarlar, gerekli bilgileri sisteme girdikten sonra bir şifre ve bir kullanıcı adı alır. Kendi şifre ve kullanıcı adını ile makale gönderme sistemine kayıt olduktan sonra yazının işleme alınmasında bir gecikme olmaması için gerekli tüm bilgileri sağlamak için sistemin yönergelerini dikkatlice okuyunuz. Makaleyi ve tüm şekil, tablo ve ek dökümanları ekleyiniz. Ayrıca üst yazı ve "Telif Hakkı ve Finansal Durum" formunu ve yazının tipine göre aşağıda belirtilen kılavuzların kontrol listesini ekleyiniz.

Makale Hazırlama Kuralları

Türk Kolon ve Rektum Hastalıkları Dergisi "Biyomedikal Dergilere Gönderilen Makaleler için Gerekli Standartları" izler. (International Committee of Medical Journal Editors: Br Med J 1988; 296: 401-5).

Yazarlar yayınlannı gönderirken, çalışmalarının türünü ve uygulanan istatistik yöntemlerini "Tıbbi Dergilere Gönderilen Makaleler için İstatistiksel Raporlama Rehberi"ne uygun olarak belirtmelidir (Bailar JC III, Mosteller F. Ann Intern Med 1988;108:266-73).

Araştırma makalesi, sistematik değerlendirme ve meta-analiz hazırlanması aşağıdaki çalışma tasarımı kurallarına uymak zorundadır; (CONSORT statement for randomized controlled trials (Moher D, Schulz KF, Altman D, for the CONSORT Group).

The CONSORT statement revised recommendations for improving the quality of reports of parallel group randomized trials. JAMA 2001; 285:1987-91) (<http://www.consort-statement.org/>);

PRISMA statement of preferred reporting items for systematic reviews and meta-analyses (Moher D, Liberati A, Tetzlaff J, Altman DG, The PRISMA Group. Preferred Reporting Items for Systematic Reviews and Meta-Analyses: The PRISMA Statement. PLoS Med 2009; 6(7): e1000097.) (<http://www.prisma-statement.org/>);

STARD checklist for the reporting of studies of diagnostic accuracy (Bossuyt PM, Reitsma JB, Bruns DE, Gatsonis CA, Glasziou PP, Irwig LM, et al., for the STARD Group. Towards complete and accurate reporting of studies of diagnostic accuracy: the STARD initiative. Ann Intern Med 2003;138:40-4) (<http://www.stard-statement.org/>);

STROBE statement, a checklist of items that should be included in reports of observational studies (<http://www.strobe-statement.org/>);

MOOSE guidelines for meta-analysis and systemic reviews of observational studies (Stroup DF, Berlin JA, Morton SC, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting Meta-analysis of observational Studies in Epidemiology (MOOSE) group. JAMA 2000; 283: 2008-12).

Metin Biçimlendirme

Yazılar Word programı ile hazırlanarak teslim edilmelidir.

- Metin için normal, düz yazı tipi kullanın (örneğin, 10 punto Times Roman).

- Sayfa numarası için otomatik sayfa numaralandırma işlevini kullanın.

Turkish Journal of COLORECTAL DISEASE



Yazarlara Bilgi

- Alan fonksiyonları kullanmayın.
- Girintiler için sekme durakları (Tab) kullanın, ara çubuğu ve diğer komutlar kullanmayın.
- Tablo yapmak için diğer işlevleri değil, elektronik tablo fonksiyonunu kullanın.
- Dosyanızı .docx formatında (Word 2007 veya üstü) ya da .doc formatında (eski Word sürüm) kaydedin.
- Giriş sayfası
- Tüm yazılar, makale türü ne olursa olsun, aşağıdakileri içeren bir başlık sayfası ile başlamalıdır:
- Makalenin başlığı;
- Makalenin kısa başlığı;
- Yazarların isimleri, isimlerinin baş harfleri ve her yazının akademik ünvanı;
- Her yazının görevi;
- Her yazının kurumu;
- Yazının adı ve e-posta adresi;
- Herhangi bir yazının olası bir çıkar çatışması olduğunu teyit eden bir ifade, aksi takdirde çatışma olmadığını belirten bir açıklama;
- Özet, kaynaklar, tablo ve şekiller hariç kelime sayısı;
- Varsa yayının yayınlanmış olduğu bilimsel toplantının tarihi, yeri ve varsa kongre özet kitabındaki özeti.

Makale Tipleri

Orijinal Makaleler

Bu kategori, klinik ve temel bilimlerde orijinal araştırmaları içerir. Yayın orijinal olmalı ve başka bir dergide yayınlanmış/gönderilmiş ya da kabul edilmiş olmamalıdır. Yazarlar, herhangi biri tarafından bir dergiye gönderilmiş, baskıda veya basılmış ilgili herhangi bir çalışmaya atıfta bulunmak istiyorlarsa açıkça atıfta bulunulmalı ve kaynak gösterilmelidir.

Tüm klinik çalışmalar, Uluslararası Tıp Dergisi Editörler Komitesince (ICMJE) kabul gören bir kayıt sistemine kayıtlı olmalıdır. Bunun için <http://www.icmje.org/faq.html> adresine müracaat edin. Randomize kontrollü çalışmaların yazarları da, www.consort-statement.org adresinden başvurulabilen CONSORT kılavuzuna uymalıdır ve yayınlarıyla birlikte CONSORT kontrol listesi ve akış diyagramı tebliğ edilmelidir. Akış şeması olarak www.consort-statement.org adresinde bulunan MS Word şablonunun kullanılması ve bunun yayının içinde bir alıntı veya bir figür olarak yerleştirilmesi gereklidir. Buna ek olarak, sunulan yayımlar her yayına spesifik verilen özel kayıt numarasını içermelidir.

Tüm yazarların, insan üzerindeki çalışmalar ve hayvan deneylerine etik standartlara uymaları beklenmektedir. İnsan üzerindeki veya laboratuvar hayvanları içeren çalışmalarda, yazarların yayının Gereç ve Yöntem kısmında deney protokolünün ilgili kurumsal inceleme komitesi tarafından onaylandığını ve sorumlu devlet kurumu kurallarına uyduğunu açık bir dille açıklamaları gereklidir. İnsan üzerindeki çalışmalarda kurumsal inceleme kurulu onayına ek olarak, aydınlatılmış onam da bulunmalıdır.

Orijinal Makaleler (özet, kaynaklar, tablolar, rakamlar hariç) 3000 kelime ve dört figürü aşmamalıdır.

Orijinal Makaleler aşağıdaki gibi organize edilmelidir:

Özet: Özet 250 kelimeyi geçmemeli ve şunları içermelidir;

Amaç: Çalışmanın amacı nedir?

Yöntem: Kullanılan yöntem ve materyaller (örneğin hayvanlar) veya hastalar ya da konu (sağlıklı gönüllüler gibi) hakkında kısa bir açıklama içermelidir.

Bulgular: Ana bulgular nelerdir?

Sonuç: Çalışmanın ana sonuçları ve etkileri nelerdir?

Anahtar kelimeler: Özeti altında en az 3 anahtar kelime veriniz. Kısaltmalar anahtar kelime olarak kullanmayınız.

Giriş: Açık bir dille çalışmanın amaç ve gerekçesini belirtin ve çalışmanın arka planını açıklarken sadece en önemli kaynaklardan alıntı yapın.

Gereç ve Yöntem: Gözlemsel veya deneysel deneklerin (hastalar, deney hayvanları veya kontrol grupları dahil) seçim şeklini açıklayın. Deney protokolünün ilgili kurumsal inceleme komitesi tarafından onaylandığını ve ilgili devlet kurumu kurallarına uyduğunu açık bir dille açıklayın. İnsan çalışması durumunda, tüm şahısların aydınlatılmış onamlarının alındığını açık bir dille belirtin. Yöntem, cihaz ve türleri tanımlayın (Parantez içinde üretici firma adı ve adresi)** Uygulanmış olan tüm prosedürler, diğer çalışmacıların aynı deneyi tekrar edebilecekleri detay ve netlikte anlatılmalıdır. İstatistiksel yöntemler de dahil olmak üzere yerleşik ve yaygın olarak bilinen çalışma yöntemleri için kaynaklar belirtilmelidir. Yayınlanmış ancak yaygın olarak bilinmeyen yöntemler için ise kaynaklar ve kısa tanımlamalar verilmelidir. Kullanma sebepleri ve limitasyonları belirtilmelidir.

Bulgular: İstatistiksel yöntemlerle desteklenmiş bulgularınızı ayrıntılı olarak sunun. Şekil ve tablolar metni tekrar değil, takviye etmelidir. Verilerin hem metinde hem figür olarak verilmemesi gerekir. Metin veya figürden birisi olarak verilmesi yeterlidir. Sadece kendi önemli izlenimlerinizi belirtin. Kendi izlenimlerinizi diğerlerinininkiyle karşılaştırmayın. Bu tür karşılaştırma ve yorumlar tartışma bölümünde yapılmalıdır.

Tartışma: Bulgularınızın önem ve anlamını vurgulayın ancak bulgular kısmında verilenleri tekrarlamayın. Fikirlerinizi yalnızca bulgularınızla kanıtlayabildiklerinizle sınırlı tutun. Bulgularınızı diğerlerinininkiyle karşılaştırmayın. Bu bölümde yeni veriler bulunmamalıdır.

Teşekkür: Sadece çalışmaya ciddi katkılarda bulunmuş kişilere teşekkür edin. Yazarlar ismen teşekkür ettikleri herkesten yazılı izin almak zorundadır. Teşekkür kısmına "Yazarlarteşekkür eder" şeklinde başlayın.

Yazarlık ve Katkı Sağlayanlar: Dergi, biyomedikal dergilere gönderilen yayınlara yönelik ICMJE tavsiyelerini izler. Buna göre "yazarlık" aşağıdaki dört kritere dayalı olmalıdır:

Yazar;

- Yayının konsept veya dizaynına, çalışmanın verilerinin elde edilmesine, analizine ve yorumlanmasına önemli katkılar veren; ve

- İşi hazırlayan veya entelektüel içerik açısından eleştirel biçimde gözden geçiren; ve

- Yayınlanacak son şekli onaylayan; ve

- Çalışmanın her bir bölümünün doğruluğu ve bütünlüğü ile ilgili sorunları uygun bir şekilde inceleleyen ve çözüm sağlayan sorumlu kişidir.

Bu şartların hepsini sağlamayan diğer tüm katılımcılar yazar değil, "Teşekkür" bölümünde anılması gereken katkı sağlamış kişilerdir.

Kaynaklar: Kaynakları 1'den başlayarak Arap rakamları ve alfabetik sıra ile verin. Kaynak numaraları cümle sonunda noktadan sonra üstte küçük rakamlar şeklinde (superscript) yazılmalıdır. Kısaltmalar için gerekli standartları <http://www.bilimterimleri.com> adresinde bulunan Türk Bilim Terimleri Kılavuzu'ndan edinin.

Dergi başlıkları "Cumulated Index Medicus" kısaltmalarına uygun olmalıdır.

Dergiden: Yazar/yazarların soyadı ve adının ilk harfi, makale başlığı, dergi başlığı ve derginin özgün kısaltması, yayın tarihi, baskı, kapsayıcı sayfa numaralarını içermelidir.

Örneğin: 1. Dilaveris P, Batchvarov V, Gialafos J, Malik M. Comparison of different methods for manual P wave duration measurement in 12-lead electrocardiograms. Pacing Clin Electrophysiol 1999;22:1532-1538.

Kitap Bölümü: Yazar/yazarların soyadı ve adının ilk harfi, bölüm başlığı, kitap editörleri, kitap başlığı, basım, yayın yeri, yayın tarihi, kapsadığı sayfa numaralarını içermelidir

Örneğin: 1. Schwartz PJ, Priori SG, Napolitano C. The Long QT Syndrome. In: Zipes DP, Jalife J, eds. Cardiac Electrophysiology. From Cell to Bedside. Philadelphia; WB Saunders Co. 2000:597-615.

Tablolar: Tüm tablolar Arapça sayılarla numaralandırılmalıdır. Tüm tablolardan metin içerisinde numara sırası ile bahsedilmelidir. Her tablo için tablonun içeriği hakkında bilgi veren bir başlık verin. Başka yayından alıntı olan tüm tablolar tablonun alt kısmında kaynak olarak belirtin. Tabloda dipnotlar tablonun altında, üst karakter olarak küçük harflerle verilmelidir. İstatistiksel anlamı değerler ve diğer önemli istatistiksel değerler yıldız ile işaretlenmelidir.

Şekiller: Şekillerin "Windows" ile açılması gerekir. Renkli şekiller veya gri tonlu görüntüler en az 300 dpi olmalıdır. Şekiller ana metinden ayrı olarak ".tiff", ".jpg" veya ".pdf" formatında kaydedilmelidir. Tüm şekil ayrı bir sayfada hazırlanmalı ve Arap rakamları ile numaralandırılmalıdır. Her şekilde kendisindeki işaret ve semboller açıklayan bir alt yazı olmalıdır. Şekil gönderme için yazardan hiçbir ek ücret alınmaz.

Ölçü Birimleri ve Kısaltmalar: Ölçü birimleri System International (SI) birimleri cinsinden olmalıdır. Kısaltmalardan başlıkta kaçınılmalıdır. Sadece standart kısaltmalar kullanın. Metinde kısaltma kullanılırsa ilk kullanıldığı yerde tanımlanmalıdır.

İzinler: Yazarlar yayınlara önceden başka bir yerde yayınlanmış şekil, tablo, ya da metin bölümleri dahil etmek isterlerse telif hakkı sahiplerinden izin alınması ve bu izin belgelerinin yayına beraber gönderilmeye gönderilmesi gerekmektedir. Böyle bir belgenin eşlik etmediği her materyalin yazara ait olduğu kabul edilecektir.

Davetli (Talep üzerine yazılan) Derlemeler

Özet uzunluğu: 250 kelimeyi aşmamalıdır.

Makale uzunluğu: 4000 kelimeyi aşmamalıdır.

Kaynak sayısı: 100 kaynağı aşmamalıdır.

Turkish Journal of COLORECTAL DISEASE



Yazarlara Bilgi

Derlemeler, üzerine konuyla ilgili yeni bir hipotez ya da çalışma oturtulabilecek bir sonuç içermelidir. Literatür taraması metodlarını veya kanıt düzeyi yöntemlerini yayınlamayın. Derleme makaleleri hazırlayacak yazarların ilgili konuda önceden araştırma makaleleri yayınlamış olması gerekir. Çalışmanın yeni ve önemli bulguları sonuç bölümünde vurgulanır ve yorumlanmalıdır. Derlemelerde maksimum iki yazar olmalıdır.

Olgu Sunumları

Özet uzunluğu: 100 kelimeyi aşmamalıdır.

Makale uzunluğu: 1000 kelimeyi aşmamalıdır.

Kaynak sayısı: 15 kaynağı aşmamalıdır.

Olgu Sunumları aşağıdaki gibi yapılandırılmalıdır:

Özet: Olguyu özetleyen bir yapılandırılmamış özet (gereç ve yöntem, bulgular, tartışma gibi bölümlerin olmadığı).

Giriş: Kısa bir giriş (tavsiye edilen uzunluk: 1-2 paragraf).

Olgu Sunumu: Bu bölümde ilk tanı ve sonuç da dahil olmak üzere olgu ayrıntılı olarak anlatılır.

Tartışma: Bu bölümde ilgili literatür kısaca gözden geçirilir ve sunulan olgunun, hastalığa bakışımızı ve yaklaşımımızı nasıl değiştirebileceği vurgulanır.

Kaynaklar: Vancouver tarzı, (yukarıda 'Kaynaklar' bölümüne bakınız).

Teşekkür

Tablolar ve şekiller

Teknik Notlar

Özet uzunluğu: 250 kelimeyi aşmamalıdır.

Makale uzunluğu: 1200 kelimeyi aşmamalıdır.

Kaynak Sayısı: 15 kaynağı aşmamalıdır.

Teknik Notlar, yeni bir cerrahi tekniğin açıklanmasını ve az sayıda olguda uygulanmasını içermektedir. Büyük bir atılım/değişikliği temsil eden bir tekniğin sunulması durumunda tek bir olgu yeterli olacaktır. Hastanın takip ve sonucu açıkça belirtilmelidir.

Teknik Notlar aşağıdaki gibi organize edilmelidir:

Özet: Aşağıdaki gibi yapılandırılmalıdır:

Amaç: Bu çalışmanın amacı nedir?

Yöntem: Kullanılan yöntemlerin, hastalar ya da sağlıklı gönüllülerin veya hayvanların tanımı, malzemeler hakkında kısa bir açıklama.

Bulgular: Ana bulgular nelerdir?

Sonuç: Bu çalışmanın ana sonuçları ve etkileri nelerdir?

Endikasyonları

Yöntem

Diğer yöntemlerle karşılaştırılması: Avantaj ve dezavantajları, zorluklar ve komplikasyonlar.

Kaynaklar: Vancouver tarzı (yukarıda 'Kaynaklar' bölümüne bakınız)

Teşekkür

Tablolar ve şekiller; alt yazıları dahil

Video Makale

Makale Uzunluğu: 500 kelimeyi aşmamalıdır.

Kaynak Sayısı: 5 kaynağı aşmamalıdır.

Tanıyı, uygulanan cerrahi tekniği ve sonucu açıklayarak olguyu kısaca özetleyiniz. Uygun şekilde adlandırılmış ve referans edilmiş video materyalleri ile tüm önemli noktaları, örneğin; yeni cerrahi tekniği, belirtiniz. Materyaller, yazarların cerrahi tekniğini anlattıkları veya karşılaştıkları ilginç vakalardan oluşmalıdır.

Teknik Gereklilikler: Veriler, makale yükleme sırasında diğer dosyalarla birlikte eklenmelidir. Video süresinin 10 dakikayı geçmemesi kaydıyla dosya boyutu maksimum 350 MB olmalı ve 'MOV, MPEG4, AVI, WMV, MPEGPS, FLV, 3GPP, WebM' formatlarından biri kullanılmalıdır. 100 MB'yi aşmayan video dokümanları sisteme yüklenebilir. Daha büyük video dokümanları için lütfen iletisim@galenos.com.tr adresinden bizimle iletişime geçiniz. Tüm video seslendirmeleri İngilizce olmalıdır. Video atıfları, Şekil veya Tablo atıfları ile aynı biçimde kullanılmalıdır. Örneğin; "...Bunu gerçekleştirmek için, yeni bir cerrahi teknik geliştirdik (Video 1)." Video materyallerinde isim ve kurumlar yer almamalıdır. Kabul edilen makalelerin video materyalleri online yayınlanacaktır.

Editöre Mektuplar

Makale uzunluğu: 500 kelimeyi aşmamalıdır.

Kaynak Sayısı: 10 kaynağı aşmamalıdır.

Türk Kolon ve Rektum Hastalıkları Dergisi'nde yayınlanan makaleler hakkında yorumlar memnuniyetle kabul edilir. Özet gerekli değildir, ancak lütfen kısa bir başlık ekleyiniz. Mektuplar bir şekil veya tablo içerebilir.

Editöryal Yorumlar

Makale uzunluğu: 1000 kelimeyi aşmamalıdır.

Kaynak Sayısı: 10 kaynağı aşmamalıdır.

Editöryal yorumlar sadece editör tarafından kaleme alınır. Editöryal yorumlarda aynı konu hakkında başka yerlerde yayınlanmış yazılar hakkında fikir veya yorumlar belirtilir. Tek bir yazar tercih edilir. Özet gerekli değildir, ancak lütfen kısa bir başlık ekleyiniz. Editöryal gönderimler revizyon/gözden geçirme talebine tabi tutulabilir. Editörler, metin stilini değiştirmeye hakkını saklı tutar.

Etik

Bu dergi, bilimsel kayıtların bütünlüğünü korumayı taahhüt etmektedir. Yayın Etik Komitesi (COPE) üyesi olarak, dergi olası olumsuz davranışlarla nasıl başa çıkılacağı konusunda Yayın Etik Komitesi (COPE) kılavuzlarını takip edecektir.

Yazarlar araştırma sonuçlarını yanlış sunmaktan; derginin güvenilirliğine, bilimsel yazarlık profesyonelliğine ve en sonunda tüm bilimsel çabalara zarar verebileceğinden dolayı, sakıncalıdır. Araştırma bütünlüğünün sürdürülmesi ve bunun sunumu, iyi bilimsel uygulama kurallarını takip ederek başarılıdır. Bu da şunları içerir:

- Yazılı eser değerlendirilmek üzere eş zamanlı birden fazla dergiye gönderilmemelidir.

- Yazılı eser daha önceki bir eserin geliştirilmesi olmadıkça, daha önce (kısmen ya da tamamen) yayınlanmamış olmalıdır. [Metnin yeniden kullanıldığı imasından kaçınmak için tekrar kullanılabilir materyallerde şeffaflık sağlayın ("self-plagiarism" kişinin kendinden intihali)].

- Tek bir çalışma; sunum miktarını arttırmak için birçok parçaya bölünmemeli ve zaman içinde aynı ya da çeşitli dergilere gönderilmemelidir. (örneğin "salam-yayınçılık" "salamizasyon").

- Veriler, sonuçlarınızı desteklemek için fabrikasyon (uydurma) ya da manüple edilmiş olmamalıdır.

- Yazarın kendine ait olmayan hiçbir veri, metin veya teori kendininmiş gibi sunulmamalıdır (intihal). Diğer eserlerin kullanımı, (eserin birebir kopyalanması, özetlenmesi ve/veya başka kelimeler kullanarak açıklanması da içeren) ya telif hakkı korunacak şekilde izin alınarak ya da tırnak işaretinde birebir kopyalanarak uygun onay ile kullanılmalıdır.

Önemli not; Türk Kolon ve Rektum Hastalıkları Dergisi intihal taramak için bir program (iThenticate) kullanmaktadır.

- Eser sunulmadan önce sorumlu makamlardan ve çalışmanın yapıldığı enstitü/kuruluşlardan-zimnen veya açıkça-onay alınmasının yanı sıra tüm yazarlardan açıkça onay alınmış olmalıdır.

- Sunulan eserde yazar olarak ismi olanların, bilimsel çalışmaya yeterince katkısı olmuş olmalıdır ve ortak mesuliyet ve sorumluluğu olmalıdır.

Bununla beraber:

- Yazarlık veya yazarların sıra değişiklikleri eserin kabulünden sonra yapılamaz

- Yazının revizyon aşamasında, yayın öncesi veya yayınlandıktan sonra yazar isim eklenmesi veya çıkarılması istemi; ciddi bir konudur ve geçerli sebepler olduğunda değerlendirilebilir. Yazar değişikliği gerekçesi; haklı gerekçeli, inandırıcı ve sadece tüm yazarların yazılı onayı alındıktan sonra; ve yeni/siliniş yazının rolü silme hakkında ikna edici ayrıntılı bir açıklama ile kabul edilebilir. Revizyon aşamasında değişiklik olması halinde, bir mektup revise edilmiş yayına eşlik etmelidir. Yayına kabul edildikten veya yayınlandıktan sonra değişiklik olması halinde, bu istek ve gerekli dokümantasyonun yamcı yoluyla editöre gönderilmesi gerekmektedir. Gerek görüldüğünde bu isteğin gerçekleşmesi için daha fazla doküman talep edilebilir. Değişikliğin kabul veya red karar dergi editörü insiyatifindedir. Bu nedenle, yayının gönderilmesi aşamasında yazar/yazarlar; gönderecekleri ilgili yazar grubunun isim doğruluğundan sorumludur.

- Yazarlardan sonuçların geçerliliğini doğrulamak amacıyla verilerin ilgili belgelerinin istenmesi halinde bu verileri göndermek için hazır bulundurulmalıdır. Bunlar, ham veri, örnekler, kayıt vb. şeklinde olabilir.

Görevi kötüye kullanma ya da suistimal şüphesi halinde dergi COPE yönergeleri izleyerek bir soruşturma yürütecektir. Soruşturmanın ardından, iddia geçerli görünüyorsa, yazara sorunu gidermek için bir fırsat verilecektir.

Usulsüzlük, şüphe seviyesinde kaldığında; dergi editörü aşağıdaki yollardan birine başvurabilir;

- Makale halen şüpheli ise, reddedilip yazara iade edilebilir.

- Makale online yayınlanmış ise; hatanın mahiyetine bağlı olarak ya yazım hatası olarak kabul edilecek ya da daha ciddi durumlarda makale geri çekilecektir.

- Hatalı yayın ve geri çekme durumlarında açıklayıcı not yayınlanır ve yazının kurumu bilgilendirilir.

Turkish Journal of COLORECTAL DISEASE



Yazarlara Bilgi

İnsan ve Hayvan Araştırmaları

İnsan Hakları Beyannamesi

İnsan katılımlı araştırmalar: 1964 Helsinki Deklarasyonu'na ve sonrasında yayımlanan iyileştirici ilkelere uygun olmalıdır ve yazarlar tarafından kurumsal ve/veya ulusal etik kurul komitelerine başvurulup onay alınmış olduğu beyan edilmelidir.

Araştırmanın 1964 Helsinki Deklarasyonu veya kıyaslanabilir standartlara göre yürütülmesi ile ilgili şüphe durumunda, yazarlar bu durumun nedenlerini açıklamak zorundadır ve bağımsız etik kurulları veya diğer değerlendirme kurulları aracılığıyla şüphelerin giderilmesi gerekmektedir.

Aşağıda belirtilen durumlar yazı içerisinde "Kaynaklar" bölümünden önce yer almalıdır:

Etik Kurul Onayı: "Çalışmada insanlara uygulanan tüm prosedürler kurumsal ve ulusal araştırma kurullarının etik standartlarına, 1964 Helsinki Deklarasyonu'na ve sonrasında yayımlanan iyileştirici ilkelere uygun olmalıdır."

Retrospektif çalışmalarda, aşağıda belirtilen cümle yer almalıdır.

"Bu tür çalışmalarda yazılı onam gerekmemektedir."

Hayvan Hakları Beyannamesi

Araştırmalarda kullanılan hayvanların refahına saygı gösterilmelidir. Hayvan deneylerinde, yazarlar hayvanların bakımında ve kullanımında uluslararası, ulusal ve/veya kurumsal olarak oluşturulmuş kılavuzlara uymalıdır ve çalışmalar için kurumdaki veya çalışmanın yapıldığı veya yürütüldüğü merkezdeki (eğer böyle bir merkez varsa) Klinik Araştırmalar Etik Kurulundan onay alınmalıdır. Deneysel hayvan çalışmalarında "Guide for the care and use of laboratory animals" <http://oacu.od.nih.gov/regs/guide/guide.pdf> doğrultusunda hayvan haklarını koruduklarını belirtmeli ve kurumlarından etik kurul onay raporu almaldır.

Hayvanlar ile yürütülen çalışmalarda, aşağıda belirtilen durumlar yazı içerisinde 'Kaynaklar' bölümünden önce yer almalıdır:

Etik Kurul Onayı: "Hayvanların bakımı ve kullanımı ile ilgili olarak uluslararası, ulusal ve/veya kurumsal olarak oluşturulmuş tüm kılavuzlara uyulmuştur."

Eğer uygun bulduysa (komitenin bulunduğu merkezde): "Hayvan çalışmalarında yapılan tüm uygulamalar kurumsal veya çalışmanın yürütüldüğü merkez tarafından belirlenmiş etik kurallara uyumludur."

Eğer makale insan ya da hayvan katılımlı bir çalışma değilse, lütfen aşağıda yer alan uygun durumlardan birini seçiniz:

"Bu makalenin yazarları insan katılımlı bir çalışma olmadığını bildirmektedir."

"Bu makalenin yazarları çalışmada hayvan kullanılmadığını bildirmektedir."

"Bu makalenin yazarları insan katılımlı veya hayvan kullanılan bir çalışma olmadığını bildirmektedir."

Bilgilendirilmiş Onam

Bütün bireyler ihlal edilemeyecek kişisel haklara sahiptir. Çalışmada yer alan bireyler, elde edilen kişisel bilgilere, çalışmada geçen görüşmelere ve elde edilen fotoğraflara ne olacağı konusunda karar verebilmeye hakkına sahiptir. Bundan dolayı, çalışmaya dahil etmeden önce yazılı bilgilendirilmiş onam alınması önemlidir. Bilimsel olarak gerekli değilse ve

katılımcılardan (veya katılımcı yetkin değilse ebeveynlerinden veya velilerinden) basılması için yazılı onam alınmadysa, katılımcılara ait detaylar (isimleri, doğum günleri, kimlik numaraları ve diğer bilgileri) tanımlayıcı bilgilerini, fotoğraflarını ve genetik profillerini içerecek şekilde yazılı formda basılmamalıdır. Tam gizlilik sağlanmasının zor olduğu durumlarda, bilgilendirilmiş onam formu şüpheyi içerecek şekilde düzenlenmelidir. Örneğin fotoğrafla katılımcıların göz kısmının maskelenmesi gizlilik açısından yeterli olmayabilir. Eğer karakteristik özellikler gizlilik açısından değiştirilirse, örneğin genetik profilede, yazar yapılan değişikliğin bilimsel olarak sorun oluşturmadığını emin olmalıdır.

Aşağıdaki ifade belirtilmelidir:

Bilgilendirilmiş Onam: "Çalışmadaki tüm katılımcılardan bilgilendirilmiş onam alınmıştır."

Eğer makalede katılımcıların tanımlayıcı bilgileri yer alacaksa, aşağıdaki ifade belirtilmelidir:

"Makalede kişisel bilgileri kullanılan tüm katılımcılardan ayrıca bilgilendirilmiş onam alınmıştır."

DEĞERLENDİRME SÜRECİ

Türk Kolon ve Rektum Hastalıkları Dergisi'ne gönderilen tüm yazılar, sisteme yüklendikten sonra ilk önce editöryal kurul tarafından derginin amaç ve hedeflerine uygunluk ve temel şartları sağlama yönünden değerlendirilecektir. Yazılar, konusunda uzman dergi hakemlerine değerlendirilmek üzere gönderilecektir. Tüm kabul edilen yazılar yayımlanmadan önce, istatistik ve İngiliz dili konusunda uzman editörler tarafından değerlendirilecektir. Sayfaların ilk gözden geçirilmesinden sonra, hakem yorumları ön karar vermek için Editör'e gönderilecektir. Bu aşamada, ilk değerlendirmede bulunanların düşüncesi doğrultusunda, yazı kabul edilebilir, reddedilebilir veya yazıda düzeltme yapılması istenebilir. İlk değerlendirme sonrasında değerli bulunan makaleler için genellikle düzeltme istenir. Düzeltilen makaleler ilk karardan sonraki 2 ay içerisinde tekrar dergiye gönderilmelidir. Süre uzatılma yardımcı editörden 2 aylık süre bitmeden en az 2 hafta önce talep edilmelidir. Türk Kolon ve Rektum Hastalıkları Dergisi tarafından, 2 aylık düzeltme süresi sona erdikten sonra, yazı kabul edilmeyecektir. Düzeltme yapılan yazılar sisteme tekrar yüklendikten sonra değerlendirilmek üzere (genellikle ilk değerlendirmeyi yapan hakeme) gönderilecektir. Sonuç olarak yayımlanma kararı verildikten sonra, baskı öncesi Teknik Editör tarafından son kez değerlendirilecektir ve iletişim kurulacak olan yazara gözden geçirme ve son düzenlemeleri yapmak üzere işaretlenmiş bir nüshası elektronik ortamda gönderilecektir.

DÜZELTME SONRASI GÖNDERİLMESİ

Revize edilmiş bir versiyonu gönderirken yazar, yorumcular tarafından ele alınan her konuyu ayrıntılı olarak açıklamalı ve nokta nokta ayrıntılı olarak "yorumlara yanıt" sunmalıdır ve ardından belgenin açıklanmış kopyası bulunmalıdır (her yorumcunun yorumu nerede bulunabilir, yazarın cevap ve satır numaraları gibi yazarın değişiklikler).

Bunun yanı sıra ana revize yazı, kabul mektubu tarihinden itibaren 30 gün içinde teslim edilmelidir. Yazının revize edilmiş versiyonunun tanınan süre içinde verilmemesi durumunda, revizyon seçeneği iptal edilebilir. Yazar(lar) ek sürenin gerekli olduğunu düşünüyorsa, ilk 30 günlük süre bitmeden, uzatmayı talep etmelidir.

İNGİLİZCE YAZIM

Tüm yazılar yayımlanmadan önce profesyonel olarak "English Language Editor" tarafından değerlendirilmektedir.

KABUL SONRASI

Tüm kabul edilen makaleler editörlerden biri tarafından teknik açıdan değerlendirilecektir. Teknik inceleme tamamlandıktan sonra, makale ilgili birime gönderilerek yaklaşık bir hafta içerisinde tamamen atıf yapılabilir "Kabul Edilmiş Makale" şeklinde online olarak yayımlanacaktır.

Telif Hakkının Devri

Yayımlayan dergiyeye (veya basım ve yayma haklarının ayrı olduğu yapılarla ayrı olarak) makalenin telif hakkının devri gerekmektedir. Telif yasaları gereği bilginin yayılması ve korunması daha güvenli olarak sağlanacaktır.

Resimler

Renkli çizimlerin yayımlanması ücretsizdir.

Basım Öncesi Son Kontrol (Proof Reading)

Amaç; dizgi kontrolünü sağlamak veya dönüştürme hatalarını fark etmek, bütünlük ve netlik açısından yazıyı, tabloları ve şekilleri kontrol etmektir. Yeni bulgu ekleme, değerlerde düzeltme, başlıkta ve yazarlarda önemli değişikliklere editör izni olmadan müsadde edilmemektedir.

Online olarak yayımlandıktan sonra yapılacak değişikliklerde, Erratum üzerinden form oluşturulup makaleye erişim sağlayacak bağlantı oluşturulması gerekmektedir.

ERKEN YAYIN

Kabul edilmiş yazının baskı için tümü hazırlanırken online olarak özet hali yayımlanır. Kabul edilen yazı kontrolden geçtikten sonra, yazarlar son düzeltmeleri yaptıktan sonra ve tüm değişiklikler yapıldıktan sonra yazı online olarak yayımlanacaktır. Bu aşamada yazıya DOI (Digital Object Identifier) numarası verilecektir. Her iki forma da www.journalagent.com/krhd adresinden ulaşılabilecektir. Kabul edilen yazının yazarları elektronik ortamdaki sayfaları çıktı olarak aldıktan sonra proofreading yapmak, tüm yazıyı, tabloları, şekilleri ve kaynakları kontrol etmeyle sorumludur. Baskıda gecikme olmaması için 48 saat içinde sayfa kontrolleri yapılmış olmalıdır.

YAZIŞMA

Tüm yazışmalar dergi editöryal kuruluna ait aşağıdaki posta adresi veya e-mail adresi ile yapılacaktır.

Adres: Latilokum Sok. Alphan İşhanı No:3 Kat:2 Mecidiyeköy-Şişli, İstanbul, Türkiye

Telefon: +90 212 356 01 75-76-77

GSM: +90 (532) 300 72 36

Faks: +90 212 356 01 78

Online makale göndermek için: www.journalagent.com/krhd

Web sayfası: www.turkishjcrd.com

E-posta: info@turkishjcrd.com

Turkish Journal of COLORECTAL DISEASE



Contents/İçindekiler

Research Articles/Özgün Makaleler

- 65 **Trephine Stoma Creation Under Local Anesthesia with Sedoanalgesia**
Sedoanaljezi Eşliğinde Lokal Anestezi Altında Trefin Stoma Uygulaması
Ali Aksu, Mehmet Buğra Bozan, Nurullah Aksoy, Ayşe Azak Bozan, Nizamettin Kutluer, Burhan Hakan Kanat, Sevim Şenol Karataş, Asude Aksoy, Abdullah Böyük; Elazığ, Şanlıurfa, Turkey
- 69 **Comparison of V-Y Flap and Limberg Flap Methods in Pilonidal Sinus Surgery**
Pilonidal Sinüs Cerrahisinde V-Y Flep ile Limberg Flep Yöntemlerinin Karşılaştırılması
Ramazan Sarı, Soner Akbaba, Rıza Haldun Gündoğdu, Mustafa Ömer Yazıcıoğlu; Muş, Ankara, Turkey
- 75 **Evaluation of Etiological Risk Factors in the Development of Adult Chronic Pilonidal Disease**
Erişkin Kronik Pilonidal Sinüs Hastalığının Oluşumunda Etiyolojik Risk Faktörlerinin Değerlendirilmesi
Adnan Kuvvetli, Süleyman Çetinkunar, Alper Parlakgümüş; Adana, Turkey
- 78 **Birth Trauma: Sphincter Injuries**
Doğum Travması: Sfinkter Yaralanması
Murat Çakır, Mehmet Aykut Yıldırım; Konya, Turkey
- 82 **An Acute Abdomen Dilemma: Epiploic Appendagitis**
Bir Akut Karın Dilemması: Epiploik Appendajit
Hakan Özdemir, Oğuzhan Sunamak, Zehra Ünal Özdemir, Ferdi Cambaztepe; İstanbul, Rize, Turkey
- 85 **The Effects of Glutamine and N-Acetylcysteine on Experimental Colitis Induced by 2,4,6-Trinitrobenzene Sulphonic Acid in Rats**
Ratlarda 2,4,6-Trinitrobenzeno Sülfonik Asitle Oluşturulan Deneysel Kolit Üzerine Glutamine ve N-Asetilsisteinin Etkisi
İsmail Cem Eray, Orçun Yalav, Kubilay Dalcı, Ahmet Rencüzoğulları, Ahmet Gökhan Sarıtaş, Ömer Alabaz, Şule Menziletoğlu Yıldız, Gülşah Seydaoğlu, Figen Doran; Adana, Turkey

Case Reports/Olgu Sunumları

- 91 **Appendix Duplication Accompanied by Acute Appendicitis**
Akut Apendisit Eşlik Ettiği Apendiks Duplikasyonu
Uğur Topal, Figen Doran, Ahmet Rencüzoğulları; Adana, Turkey
- 94 **Idiopathic Nonobstructive Colo-colonic Intussusception: A Rare Clinical Presentation**
İdiyopatik Nonobstrüktif Kolo-kolonik İntussusepsiyon: Nadir Bir Klinik Tablo
Fatma Ayça Gültekin, Ramazan Kozan, Yücel Üstündağ; Zonguldak, Turkey
- 98 **Penetrating Sharp Object Injury in the Gluteal Region and Small Bowel Perforation Due to Pelvic Penetration: A Case Report**
Gluteal Bölgede Kesici Delici Alet Yaralanması: Pelvis Nafiziyeti Sonucu İnce Barsak Perforasyonu: Olgu Sunumu
Ergün Yüksel, Mehmet Akif Üstüner, Ahmet Karayığit, Lütfi Doğan; Ankara, Turkey
- 101 **Midgut Nonrotation in an Adult Patient and Ladd's Procedure**
Yetişkin Hastada Midgut Nonrotasyonu ve Ladd Prosedürü
Huriye Hande Aydınlı, Özgen Işık, Tuncay Yılmazlar; New York, United States, Bursa, Turkey

Turkish Journal of COLORECTAL DISEASE



Editorial/Editöryal

From the editors,

Dear colleagues,

We leave behind a busy working season, and we are delighted to be in the presence of you with a new issue of Turkish Journal of Colorectal Disease.

As you know, Turkish Society of Colon and Rectal Surgery has organized a very successful congress last April. The congress was attended by the honoral surgeons of Colorectal Surgery attracted great attention by national and international fields. Besides, "Masterclass" which was collaborated by Turkish Society of Colon and Rectal Surgery and European Society of Coloproctology has drawn interest widely.

My Dear Colleagues, since May, the agreement with "plexus" ended which supporting the infrastructure of Turkish Journal of Colorectal Disease. As from this issue, Turkish Journal of Colorectal Disease has now started to accept the article submission and evaluation process through its new agreement with "Manuscript Manager". Therefore, my colleagues who want to submit an article to the journal should become a member one time only again. I would like to point out that this issue should be considered with all my respect.

Two article about 'Pilonidal Sinus disease' was presented in this issue which was a lot of Authors from Turkey providing many contribution to the Medical literature about this illness. In Addition, the article presenting new local anesthetic technique for cration stoma will be worth to take attention. also, I believe that the article sharing experiences of sphincter injuries due to birth trauma and cases of acute appendagitis will draw attention.

This issue also included an experimental study investigating the effect of glutamine on the experimental colitis model. In addition, four interesting case reports were included. I believe you will read with pleasure. I wish you all a healthy summer holiday and wish to meet you in September issue...

Tahsin Çolak, MD

Surgery Professor and Editor-in-Chief

Editörden,

Değerli meslektaşlarım,

Yoğun bir çalışma sezonunu geride bırakıp, yavaş yavaş yaz aylarına girdiğimiz bu günlerde yeni ve dopdolu bir sayıyla karşınızda olmaktan mutluyuz.

Bildiğiniz gibi, Türk Kolon ve Rektum Cerrahisi Derneği Nisan ayında oldukça başarılı bir kongre düzenledi. Kolorektal cerrahinin duayen isimlerinin katıldığı bu kongre ulusal ve uluslararası alanda geniş ilgi gördü. Ayrıca ESCP (Avrupa Koloproktoloji Derneği) ve Türk Kolon ve Rektum Cerrahisi Derneği'nin ortak düzenlediği "Masterclass" da geniş ilgi gördü.

Değerli meslektaşlarım, mayıs ayı itibarıyla Türk Kolon ve Rektum Hastalıkları Dergisi alt yapısını destekleyen "pleksus" ile anlaşmasını sonlandırdı. Bu sayı itibarıyla Türk Kolon ve Rektum Hastalıkları Dergisi artık makale gönderim ve değerlendirme işlemlerini yeni anlaşma yaptığı "Manuscript Manager" üzerinden kabul etmeye başladı. Bu nedenle dergiye makale göndermek isteyen meslektaşlarımızın bir kereye mahsus yeniden üye olmaları gerekmektedir. Bu hususa dikkat edilmesini saygıyla istirham ederim.

Bu sayıda, ülkemizde sık görülen ve Türkiye'de birçok müellifin tıbbi literatüre katkı sağladığı "pilonidal sinüs hastalığı" hakkında iki makaleye yer verildi. Bununla birlikte lokal anestezi altında stoma trefini açılması ile ilgili bir diğer makalenin de dikkatinizi çekeceğini ümit ediyorum.

Doğum travmasına bağlı gelişen sfinkter yaralanmaları ve akut appendagitis olgu deneyimlerin paylaşıldığı makalelerinin dikkat çekeceğine inanıyorum.

Bu sayıda ayrıca experimental kolit modelinde glutaminin etkisinin araştıran bir deneysel çalışmaya da yer verildi.

Ayrıca birbirinden ilginç dört olgu sunumuna da yer verildi. Keyifle okuyacağınıza inanıyorum.

Hepinize sağlıklı bir yaz tatili dilerken, eylül sayısında buluşmak dileğiyle...

Prof Dr Tahsin Çolak

Cerrah Profesörü ve Baş-Editör



Trephine Stoma Creation Under Local Anesthesia with Sedoanalgesia

Sedoanaljezi Eşliğinde Lokal Anestezi Altında Trefin Stoma Uygulaması

© Ali Aksu¹, © Mehmet Buğra Bozan¹, © Nurullah Aksoy², © Ayşe Azak Bozan³, © Nizamettin Kutluer¹,
© Burhan Hakan Kanat¹, © Sevim Şenol Karataş³, © Asude Aksoy⁴, © Abdullah Büyük¹

¹University of Health Sciences, Elazığ Training and Research Hospital, Clinic of General Surgery, Elazığ, Turkey

²Şanlıurfa Siverek State Hospital, Clinic of General Surgery, Şanlıurfa, Turkey

³University of Health Sciences, Elazığ Training and Research Hospital, Clinic of Anesthesiology and Reanimation, Elazığ, Turkey

⁴Fırat University Faculty of Medicine, Department of Oncology, Elazığ, Turkey

ABSTRACT

Aim: In this paper, we aimed to evaluate the patients who underwent trephine stoma with sedoanalgesia in addition to local anesthesia and to share the results.

Method: The patients who underwent trephine stoma with sedoanalgesia in addition to local anesthesia for fecal diversion in our general surgery clinic between June 2012 and June 2017 were evaluated retrospectively. The patients were evaluated in terms of demographic characteristics, diagnosis of primary disease and complications.

Results: A total of 11 patients (F/M: 4/7) were evaluated. The mean age was 54 (± 4) years. The mean follow-up period was 18 months (range 2-30). Primary disease was rectal cancer in seven patients (63.6%), Fournier's disease in two patients (18.2%), genital cancer in one patient (9.1%), and rectovaginal fistula in one patient (9.1%). Only one patient underwent transverse colostomy and 10 patients underwent sigmoid colostomy. There were no complications related to surgical procedures. One patient died on the 14th postoperative day because of complications secondary to metastasis.

Conclusion: Trephine stoma technique, which does not require laparotomy, is a fast, reliable, and easy to use method especially in high-risk patients with poor general condition.

Keywords: Colostomy, emergent surgery, local anesthesia, minimally invasive surgery, trephine stoma

ÖZ

Amaç: Bu yazıda lokal anesteziye ilave olarak sedoanaljezi ile trefin stoma uyguladığımız hastaları değerlendirmeyi ve sonuçlarımızı paylaşmayı amaçladık.

Yöntem: Haziran 2012-Haziran 2017 tarihleri arasında genel cerrahi kliniğimizde fekal diversiyon amacıyla sedoanaljezi yardımlı lokal anestezi eşliğinde trefin stoma uygulaması yapılan hastalar retrospektif olarak incelendi. Hastalar demografik özellikleri, primer hastalığın tanısı ve komplikasyonlar açısından değerlendirildi.

Bulgular: Toplam 11 (K/E: 4/7) hastanın verileri değerlendirildi. Ortalama yaş 54 (± 4) yıl idi. Ortalama takip süresi 18 ay (2-30 arasında) idi. Hastaların primer hastalıkları 7 hastada rektum kanseri (%63,6), 2 hastada Fournier hastalığı (%18,2), 1 hastada genital kanser (%9,1), 1 hastada rektovajinal fistül (%9,1) idi. Yalnızca 1 hastaya tranvers kolostomi, 10 hastaya ise sigmoid kolostomi uygulandı. Girişimsel işleme bağlı herhangi bir komplikasyon gelişmedi. Bir hasta postoperatif 14. günde metastaza sekonder komplikasyonlar nedeniyle öldü.

Sonuç: Özellikle genel durumu iyi olmayan, yüksek riskli hastalarda, laparotomi gerektirmeyen trefin stoma tekniği hızlı, güvenilir, kolay uygulanabilen bir yöntemdir.

Anahtar Kelimeler: Kolostomi, acil cerrahi, lokal anestezi, minimal invaziv cerrahi, trefin stoma

Address for Correspondence/Yazışma Adresi: Mehmet Buğra Bozan MD,

University of Health Sciences, Elazığ Training and Research Hospital, Clinic of General Surgery, Elazığ, Turkey

Phone: +90 530 341 50 44 E-mail: bbozan@yahoo.com ORCID ID: orcid.org/0000-0001-5573-2645

Received/Geliş Tarihi: 22.10.2018 Accepted/Kabul Tarihi: 28.10.2018



Introduction

In colorectal surgery, stoma applications are the process of opening the intestinal tract to the abdominal wall. The history of surgical stoma dates back 200 years.¹ Stomas can be either temporary or permanent and their indications are very various. In colorectal surgery, the stoma is mostly performed for incontinence requiring fecal diversion, Fournier's gangrene, complex anorectal fistulas, anorectal trauma, and obstructed anorectal tumors. The techniques for the creation of stomas for fecal diversion may vary depending on the indication and the condition of the patient.² The most important determinants for the selection of the type of stoma are the location of the stoma, the technique used, and the characteristics of the patient.¹ This procedure is usually done by laparotomy, but it can also be performed without a laparotomy.

Trepine stoma technique is a less invasive procedure without laparotomy. Trepine as a medical term means the removal of a tissue by circular incision. In this technique, a circular incision is made to the previously determined area in the abdominal wall and a stoma is created with the intestines brought from the abdomen. It is a method that can be preferred especially in patients with poor general condition, comorbidity and complication.

In the literature, there are some articles related to this technique and the procedure was performed under general anesthesia or regional anesthesia in most of them. In our clinic, the application was performed with sedoanalgesia in addition to local anesthesia. In this paper, we aimed to evaluate the patients who underwent trephine stoma with sedoanalgesia in addition to local anesthesia and to share the results.

Materials and Methods

The patients who underwent trephine stoma with sedoanalgesia in addition to local anesthesia for fecal diversion in our general surgery clinic between June 2012 and June 2017 were retrospectively evaluated. Data of the patients were obtained from personal records and computer records. Patients with missing data were excluded from the study. A total of 11 patients were included in the study. The patients were evaluated in terms of demographic characteristics, diagnosis of primary disease and complications. All patients were consulted with the anesthesia clinic as routine procedure and the risks of anesthesia were determined.

Statistical Analysis

Statistical analysis of the study was performed using SPSS 20 (SPSS for Windows, Inc., Chicago, IL, USA). Data were presented as percentage and arithmetic mean \pm standard deviation (minimum-maximum).

Technique

The procedure was performed under sterile conditions in the operating room. Colostomy area was marked in the left lower quadrant while the patients were sitting, standing and lying. After a single dose of antibiotic prophylaxis, the patient was draped and prepped as if laparotomy was to be performed. All patients received nasal oxygen at 2 liters/min during the procedure. All medications were administered intravenously.

Initially, all patients were administered midazolam 0.05 mg/kg and fentanyl 1 mcg/kg by anesthesiologist. Ramsay sedation scale that was developed by M. A. Ramsay in the mid-1970s was used to determine the level of sedation of patients.³ Ramsay score of the patients was kept between 3-4. This scale consists of a total of six items, each of which contains three levels of awake and asleep.⁴ These are as follows: patient awake, anxious, agitated, or restless; patient awake, cooperative, orientated, and tranquil; patient drowsy, with response to commands; patient asleep, brisk response; patient asleep, sluggish response and patient has no response. The first three responses are evaluated in the awake patient, and the other three responses are evaluated by glabella tap or loud auditory stimulus. Midazolam 0.03 mg/kg and fentanyl 0.5 mcg/kg IV were administered when additional dose was needed.

While the patient was sedated, rectus muscle and sheath were infiltrated with lidocaine or bupivacaine. Lidocaine is faster, but short-acting, while bupivacaine is slower and much longer effective. The dose limitation for both is independent of each other. The recommended dose by the Food and Administration is 7 mg/kg for lidocaine. The maximum reported dose range is 200-300 mg. The maximum dose for bupivacaine is 175 mg. If adrenaline is added, these can be increased up to 500 and 225 milligrams.⁵

Local anesthesia including intradermal, subdermal and deep subcutaneous is performed in the previously marked area. An elliptical incision in normal colostomy size is used to pass the skin and subcutaneous fascias to reach the fascia of the left rectus muscle. Next, local anesthetic agent infiltration is performed on the lateral of the rectus abdominis muscle toward the aponeurosis of the internal, external oblique and transversus abdominis muscle. After the fascia is incised, the muscles are opened and the peritoneum is reached. At this stage, the intestinal loop is taken out of the opening by Babcock clamp (loop colostomy) and the baguette is passed. The colostomy mouth is then sutured (Figure 1).

Results

The data of 11 patients included in the study were evaluated. The mean age of the patients was 54 ± 4 years (range: 37-81).

Four of the patients were female (36%) and seven were male (64%). Regarding primary diagnosis, seven patients had rectal cancer (63.6%), two patients had Fournier's disease (18.2%), one patient had genital cancer (9.1%), and one patient had rectovaginal fistula (9.1%). Six patients (54.5%) had comorbidities, including diabetes mellitus, ischemic heart disease, chronic obstructive pulmonary disease, and other organ metastases due to the rectum tumor (lung, liver, adrenal). Only one patient underwent transverse colostomy and 10 patients underwent sigmoid colostomy. Five of the patients with rectal cancer had complete obstruction and 2 patients had partial obstruction. These 7 patients were



Figure 1. The patient who underwent trephine stoma

treated under emergency conditions and the other patients were treated electively. No complication was observed in any patient due to interventional procedures. One patient died on the 14th postoperative day because of complications secondary to metastasis. The mean follow-up period was 18 months (range: 2-30) (Table 1).

No anesthetic drug-related complication was observed during or after the procedure. Wound site infection developed in one patient who had ostomy due to Fournier's gangrene. In all patients, oral intake was started on the 1st postoperative day. No postoperative ileus was observed in any patient. Three of the patients who underwent ostomy due to rectal cancer were operated laparoscopically during the follow-up period.

Discussion

Stoma is a Greek word meaning "opening" or "mouth". Intestinal stoma means that the intestine is mouthed on the abdominal wall. It is divided into two, as permanent or temporary. Furthermore, it is classified as end ileostomy (Brooke ileostomy), loop ileostomy, continent (Kock) ileostomies, loop colostomy, double-barrel colostomy (Mikulicz), Hartmann's procedure, and permanent end colostomy according to opening type.⁶

Stomas can be opened with laparotomy or laparoscopy or without laparotomy. Trepine stoma technique, which is opened without laparotomy, was first described by Senapati. Trepine stoma technique can be performed as ileostomy or colostomy.^{2,7} Trepine stoma technique, which does not require laparotomy in high-risk patients, is a fast, reliable and easy to use method. Previous abdominal surgery is

Table 1. The characteristics of the patients

Patient ID	Age	Gender	The underlying reason	Comorbidities	Colon segment
1	74	M	Fournier's gangrene	COPD	Transverstomy
2	68	M	Obstructed rectum cancer	Ischemic heart disease	Left colon
3	63	F	Rectovaginal fistula	Ischemic cerebrovascular disease	Left colon
4	81	F	Vulva cancer Anal invasion	Diabetes mellitus Lung metastasis	Left colon
5	37	F	Obstructed rectum cancer	No	Left colon
6	70	M	Obstructed rectum cancer	Thrombocytopenia Liver, lung and surrenal metastasis	Left colon
7	59	M	Obstructed rectum cancer	No	Left colon
8	51	F	Obstructed rectum cancer	No	Left colon
9	47	M	Fournier's gangrene	Diabetes mellitus	Left colon
10	51	M	Partial obstructed rectum cancer	No	Sigmoid colon
11	59	M	Partial obstructed rectum cancer	No	Sigmoid colon

M: Male, F: Female, COPD: Chronic obstructive pulmonary disease

usually not a contraindication. The operation time is shorter and the postoperative opiate requirement is less than the stoma performed by laparotomy.⁶ Trefin stoma is a simple, fast and the least invasive technique among the techniques of stoma creation. Because there is no separate laparotomy incision in the abdomen, postoperative recovery is faster and pain is minimal.⁷ However, it may rarely be difficult to determine the proximal and distal part of the intestine. We encountered this difficulty in our patient with genital cancer and colonoscopy was performed to determine the parts during surgery.⁸ Another topic of discussion is the length of the incision when creating the trephine stoma. In a study, it is stated that this length should be two-thirds of the stoma.⁹ The length of our incisions was the same as stoma length.

In a study on trephine stoma performed in cases with sigmoid volvulus, sigmoid colon was taken out of the body by a small incision under general anesthesia and resection was performed. Then, trephine stoma was performed.¹⁰ The most important problem of this study is the need for general anesthesia. In our study, which had small number of patients, none of the patients had sigmoid volvulus.

In the literature, there are many articles about the trephine stoma technique and most of them were performed under general anesthesia.^{11,12,13} Also, laparoscopic assisted trephine stoma technique was performed.¹³ The most important difference of our study from other studies was that it was performed under sedoanalgesia in addition to local anesthesia.

Conclusion

In patients with a high risk of anesthesia, trephine stoma creation under sedoanalgesia does not cause an additional morbidity burden due to the lack of endotracheal intubation. Also, no need for laparotomy causes less intra-abdominal adhesion in patients with rectal cancer who are candidates for neoadjuvant chemotherapy and facilitates subsequent laparoscopic surgery. Therefore, this method should be kept in mind in patients with rectum cancer or anorectal diseases requiring diversion and should be performed if necessary.

Ethics

Ethics Committee Approval: Retrospective study.

Informed Consent: Retrospective study.

Peer-review: Externally peer-reviewed.

Authorship Contributions

Surgical and Medical Practices: A.A., B.H.K., M.B.B., Concept: B.H.K., M.B.B., N.K., A.B., Design: N.K., A.A.B., N.A., A.B., Data Collection or Processing: A.A.B., N.K., S.Ş.K., A.Aksoy., Analysis or Interpretation: M.B.B., B.H.K., A.A.B., A.A., Literature Search: A.Aksoy., S.Ş.K., N.A., B.H.K., M.B.B., Writing: B.H.K., M.B.B., A.A., N.K., A.A.B.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study received no financial support.

References

1. Kapan M, Girgin S, Keleş C, Önder A, Gümüş M. Kolorektal cerrahide ostomi uygulamaları. *Türkiye Klinikleri J Gastroenterohepatoloji* 2010;17:94-106.
2. Oğuz A, Türkoğlu A, Gümüş M, Bozdağ Z, Bahadır MV, Teke F, et al. Trepine stoma: Outcomes in a single center. *J Clin Exp Invest* 2015;6:87-90.
3. Detriche O, Berre J, Massaut J, Vincent LJ. The Brussels sedation scale: use of a simple clinical sedation in patients undergoing mechanical ventilation in the intensive care. *Br J Anaesth* 1999;83:698-701.
4. Şahinoğlu AH. Yoğun bakımın doğuşu, gelişmesi ve genel ilkeler. İn Yoğun Bakım Sorunları ve Tedavi İlkeleri. Ed. 1. Baskı, Ankara, Türkiye Klinikleri Yayınevi. 1992:3-4.
5. Kulaçoğlu H. Kasık fitıklarının cerrahi tedavisinde adım-adım lokal anestezi tekniği "Teknik Not". *Turk J Surg* 2012;28:164-168.
6. Akgün EZ, Yoldaş T. İntestinal Stoma. *Kolon Rektum Hast Derg* 2012;22:133-146.
7. Neşşar G, Kayaalp C, Bil D, Karadağ S. Trefin İleostomi Tecrübemiz. *Turk J Surg* 2011;27:174-175.
8. MacCormack BJ, McCallion WA. Colonoscopic assisted trephine stoma in children with chronic constipation. *Ann R Coll Surg Engl* 2014;96:387-388.
9. Nguyen MH, Pittas F. How large should a skin trephine be for an end stoma? *Aust N Z J Surg* 1999;69:675-676.
10. Caruso DM, Kassir AA, Robles RA, Gregory MW, Tsujimura RB, Cheung P et al. Use of trephine stoma in sigmoid volvulus. *Dis Colon Rectum* 1996;39:1222-1226.
11. Hellinger MD, Al Haddad A. Minimally invasive stomas. *Clin Colon Rectal Surg* 2008;21:53-61.
12. Nylund G, Oresland T, Hultén L. The trephine stoma: formation of a stoma without laparotomy. *Eur J Surg* 1997 Aug;163:627-629.
13. Kini SU, Perston Y, Radcliffe AG. Laparoscopically assisted trephine stoma formation. *Surg Laparosc Endosc* 1996;6:371-4.



Comparison of V-Y Flap and Limberg Flap Methods in Pilonidal Sinus Surgery

Pilonidal Sinüs Cerrahisinde V-Y Flep ile Limberg Flep Yöntemlerinin Karşılaştırılması

© Ramazan Sarı¹, © Soner Akbaba², © Rıza Haldun Gündoğdu², © Mustafa Ömer Yazıcıoğlu²

¹Muş Malazgirt State Hospital, Clinic of General Surgery, Muş, Turkey

²Ankara Atatürk Training and Research Hospital, Clinic of General Surgery, Ankara, Turkey

ABSTRACT

Aim: The aim of this study was to compare the advantages of V-Y flap and Limberg flap techniques in the surgical management of complicated pilonidal sinus disease requiring extensive excision, and to review the superiorities of each other in the light of the literature.

Method: The study included 100 patients who were hospitalized with the diagnosis of pilonidal sinus disease and who were planned to undergo flap reconstruction due to the diameter of the defect. Patients were randomly divided into two groups as A and B. Fifty patients in group A underwent V-Y flap and 50 patients in group B underwent Limberg flap. Both groups were prospectively followed-up in terms of complications, recovery times, workforce losses, patient satisfaction and 2-year recurrence rates, and the findings were compared statistically.

Results: Demographic characteristics, complaints and clinical findings of both groups were similar. Although postoperative wound infection rates were similar (26% vs. 28%; $p>0.05$), wound dehiscence was more common in the Limberg group (36% vs. 26%; $p<0.05$). The wound healing process and workforce losses were longer in the Limberg flap group, but the differences were not statistically significant. There were no differences between the groups in terms of patient satisfaction and recurrence in the 2-year follow-up period.

Conclusion: The long-term recurrence rates of V-Y flap and Limberg flap techniques are similar. Wound dehiscence is significantly less in V-Y flap technique. According to the results of this study, V-Y flap method following extensive tissue loss due to excision can be considered as a good alternative.

Keywords: Pilonidal sinus, Limberg flap, V-Y flap

ÖZ

Amaç: Komplike, geniş eksizyon yapılmasını gerektirecek pilonidal sinüs hastalığının cerrahi tedavisinde popüleritesini koruyan iki yöntem olan V-Y ile Limberg flep tekniklerini karşılaştırılarak, birbirine olan üstünlüklerinin literatür ışığında gözden geçirilmesi amacıyla bir çalışma planlandı.

Yöntem: Çalışmaya pilonidal sinüs hastalığı tanısı ile yatırılan ve oluşacak defekt genişliği sebebiyle flep uygulanması planlanan 100 hasta dahil edildi. Randomize olarak 2 gruba ayrılan hastalardan A grubundaki 50 hastaya V-Y flep, B grubundaki 50 hastaya ise Limberg flep uygulandı. Her iki grup gelişen komplikasyonlar, iyileşme süreleri, iş gücü kayıpları, hasta memnuniyetleri ve 2 yıllık nüks oranları açısından prospektif olarak takip edildi ve elde edilen bulgular istatistiksel olarak karşılaştırıldı.

Bulgular: Her iki grubun demografik özellikleri, yakınmaları ve klinik bulguları benzerdi. Postoperatif yara enfeksiyonu oranları yakın olmakla birlikte (%26, %28; $p>0,005$) yara ayrışması Limberg grubunda daha fazla görüldü (%26, %36; $p<0,005$). Yara iyileşme süreci, iş gücü kayıpları Limberg flep grubunda daha uzun olmakla birlikte sonuçlar istatistiksel açıdan anlamlı değildi. Hasta memnuniyetleri ve takip edilen 2 yıllık dönemdeki nüksler açısından gruplar arasında fark yoktu.

Sonuç: V-Y flep ve Limberg flep tekniklerinin uzun dönem nüks oranları yakındır. Yara açılması V-Y flep tekniğinde anlamlı olarak daha azdır. Bu çalışmanın sonuçlarına göre, geniş doku kaybına neden olan eksizyon sonrası V-Y flep yöntemi iyi bir alternatif olarak kabul edilebilir.

Anahtar Kelimeler: Pilonidal sinüs, Limberg flep, V-Y flep



Presented in: The abstract of this study was presented as an electronic poster (EPS-0517) at the 21st National Surgery Congress held in Antalya between 11-15 April 2018 and its sample was added.

Address for Correspondence/Yazışma Adresi: Ramazan Sarı MD,

Muş Malazgirt State Hospital, Clinic of General Surgery, Muş, Turkey

Phone: +90 505 629 46 62 E-mail: sariramazan71@gmail.com ORCID ID: orcid.org/0000-0003-3492-9953

Received/Geliş Tarihi: 24.09.2018 Accepted/Kabul Tarihi: 19.11.2018

Introduction

Pilonidal sinus disease is a subcutaneous tissue disease that is common in the community and that mostly affects young adult patients. The incidence is 26/100000 in the general population and its frequency is higher in young patients and males.¹ Patients usually present with signs of abscess or sinus with chronic discharge. The treatment of the disease is important because of the painful and frequently recurring symptoms that affect the patient's lifestyle. The ideal treatment should be able to provide conditions such as quick recovery, no need for general anesthesia, outpatient treatment, minimal discomfort, minimal loss of workforce, better patient satisfaction with good cosmetic results, less recurrence and less cost. However, although many conservative and surgical methods have been defined, recurrence rates are still high and the search for the ideal treatment is ongoing.^{2,3}

Although the treatment of pilonidal sinus disease in the form of acute abscess is standard (drainage), the treatment of chronic and recurrent cases is not clear. Conservative treatment methods are recommended for simple cases with single orifice and short tracts. Surgical treatment with excision is generally accepted in cases with chronic course, multiple orifices or recurrence; however, the method of closure of the defect is controversial. In particular, there is no clear consensus on the method to be chosen for the closure of the defect in complicated or recurrent cases where the tissue loss is large.

This prospective randomized study aimed to compare popular Limberg flap and V-Y flap techniques in order to add a comment to this subject.

Materials and Methods

After the approval of the Ethics Committee of Medical Faculty of Yıldırım Beyazıt University, the patients who were admitted to General Surgery Clinic of Ankara Atatürk Training and Research Hospital between February 2012 and 2016 and who would need flap for the defect in the sacrococcygeal region after pilonidal sinus excision were included in the study. Patients who had a systemic disease that could adversely affect wound healing and who refused to participate in the study during the informed consent were excluded. When the power analysis was performed for statistical evaluation, it was calculated that 100 patients would be sufficient. Patients were labeled in groups A and B according to the order of admission. After excision of the diseased part, V-Y flap was applied to the patients in group A and Limberg flap technique was used in group B. When both groups reached 50 patients, the study was finalized.

Age, gender, duration of complaints, physical examination findings, body mass index, anesthesia method, length of

hospital stay, orifice number and length of the sinus tract were recorded. All postoperative data regarding wound infection, wound dehiscence, suture removal day, postoperative loss of sensation, time to return to work, recurrence and its treatment, patient satisfaction and cosmetic concerns were recorded.

The surgical field of all patients was shaved on the operating table just before the operation. Povidone-iodine was used for skin cleansing. All patients underwent antibiotic prophylaxis with 1 g of cefazolin sodium during anesthesia induction. After the patients were taken to the operating table in the prone jackknife position, the excision area and the area suitable for the flap technique were marked with a pen.

In group A, the tissue was totally excised down to presacral fascia with a rectangular incision including all sinuses and pits until the presacral fascia was visualized. Following hemostasis, a V shaped incision was used to incise the right gluteal region up to gluteus muscle fascia and the tissue to be used for the flap was mobilized. The obtained fasci-cutaneous flap was shifted to the opposite side in Y shape and was sutured to presacral fascia and subcutaneous skin with 1/0 polyglactine sutures. The skin was closed in interrupted 2/0 monofilament polypropylene sutures (Figure 1). A hemovac drain was used in all patients.

In group B, the tissue was totally excised down to presacral fascia with a rhomboid incision including all sinuses and pits. Following hemostasis, the flap from the right gluteal region was raised so that it included skin, subcutaneous fat, and the fascia overlying gluteus maximus, and rotated to cover the defect. Rhomboid was inserted into the defect so that the lower end did not remain in the intergluteal groove. The flap was sutured to presacral fascia and subcutaneous skin with 1/0 polyglactine sutures. The skin was closed in interrupted 2/0 monofilament polypropylene sutures. A hemovac drain was used in all patients (Figure 2).

All surgeries were performed by the same team. The wound was cared with pressure dressings and the drains were preserved until the daily drainage was below 10 mL. The wound infection was defined as presence of symptoms such as erythema, pain, redness, and induration, or surgeon opening the wound or defining the wound as infected, as well as purulent drainage not exceeding the skin and subcutaneous tissue in the wound within 30 days after the surgery. Five days of oral ampicillin + sulbactam treatment was performed in patients with wound infection. Spontaneous dehiscence without any evidence of infection in the first month was defined as wound dehiscence. The sutures of the patients in both groups were removed in our outpatient clinic and the day of removal was recorded in the

follow-up form. Patients in follow-up were checked in our outpatient clinic at the 6th, 6th and 24th months. The patients who could not come to the hospital were interviewed by telephone in terms of their time to return to work, operation satisfaction and recurrence, and their results were recorded. While evaluating the patient satisfaction levels, the patients were asked to score with a number between 1 and 5 (highest satisfaction 5, least satisfaction 1).

Statistical Analysis

SPSS for Windows 20.0 package program (SPSS Inc, Chicago, Illinois, USA) was used for statistical analysis of the data. Continuous variables were expressed as mean ± standard deviation. Continuous variables were compared using t-test and categorical variables were compared using chi-square test. ANOVA test and post hoc Tukey test were used for the comparison of parametric data between the groups. Kruskal-Wallis and chi-square test were used for comparison of nonparametric values and percentages. A p value of <0.05 was considered statistically significant in all tests.



Figure 1. V-Y flap method



Figure 2. Limberg flap method

Results

Ninety-three patients were male and seven were female, and the mean age was 26.8 years (range, 15-46). There was no difference between the two groups in terms of demographic characteristics and the anatomical structure of the disease ($p>0.05$) (Table 1).

Of the patients in group A, the number of patients with previous surgeries were as follows: three had primary repair, one had Limberg flap, one had Karydakis flap, one had marsupialization and five had only drainage. Of five patients with previous surgery in group B, three had primary repair, one had V-Y flap, and one had only drainage. There was no significant difference between the groups in terms of previous surgery data ($p>0.05$).

There were no statistically significant differences between the two groups in terms of preoperative symptoms, orifice number, sinus tract length, type of anesthesia used for surgery and length of hospital stay. Complaints of all patients were pain, swelling and discharge in the sacral region. In all cases, there was more than one sinus orifice and there was also sinus orifice outside the midline. In other words, there was a need for an excision that would require a flap in all patients.

There were no differences between the groups in terms of postoperative complications, length of hospital stay, time of suture removal, loss of sensation, and time to return to work (Table 2). Wound infections were found in 13 patients in group A on day 7 (range, 4-13) and in 14 patients in group B on day 7 (range, 3-13). Wound dehiscence was observed on day 6 (range, 3-10) in

Table 1. Characteristics of groups

	V-Y group (A) (n=50)	Limberg group (B) (n=50)	p
Age, years	26.1 (15-45)	27.66 (15-46)	0.327
Gender, M/F	43/7	50/0	0.075
BMI, kg/m ²	24.93 (20.21-32.41)	25.95 (22.49-34.61)	0.189
Comorbidities	0	1	0.317
Previous surgery	11	5	0.103
Duration of complaints, month	18.34 (1-96)	17.27 (0.5-120)	0.418
Number of sinus orifices	3 (2-5)	3 (2-6)	1.00
Tract length, cm	3.03 (1-7)	2.7 (1-7)	0.306

M: Male, F: Female, BMI: Body mass index


group A and on day 9 (range, 3-14) in group B. While there was no difference between the groups in terms of time of dehiscence, it was seen in more patients in Limberg group (36% vs. 26%; $p < 0.05$).

Recurrence was observed in two patients (mean: 19 months) in group A and in three patients in group B (mean: 11 months) in the first 2 years. Although there was no significant difference between the groups in terms of recurrence, it was observed that recurrences developed earlier in the Limberg group. Contrary to the predicted, there was no difference in recurrence rates between obese patients and non-obese patients, and patients with and without infection ($p > 0.05$). Although it was observed that patients in group A were more satisfied with cosmetic results, there was no statistically significant difference between the two groups in terms of overall satisfaction ($p > 0.05$) (Graphic 1).


Table 2. Postoperative findings

	V-Y group (A)	Limberg group (B)	P
Wound infection	13	14	0.823
Wound dehiscence	13	18	0.004
Length of hospital stay, day	1.84 (1-5)	2.02 (1-7)	0.564
Suture removal, day	17.8 (12-30)	18.7 (12-30)	0.354
Loss of sensation	5	7	0.540
Return to work, day	27.86 (7-62)	27.38 (15-60)	0.832
Patient Satisfaction	4.28 (1-5)	4.14 (2-5)	0.499
Recurrence	2	3	0.648

Evaluation of patient satisfaction in the V-Y group



Evaluation of patient satisfaction in the Limberg group



Graphic 1. Evaluation of patient satisfaction

Discussion

Despite the definition of many methods for the closure of the wound in complicated pilonidal sinus patients treated with surgical excision that leaves a wide tissue defect, there is no consensus on optimal flap technique. The main objective is to ensure that the patient is operated with a technique with a low morbidity and mortality rate, and that he/she returns to work as soon as possible. For this, the complication rate of the procedure should be low, the hospital stay should be short and the recurrence rates should be low.^{2,4,5} As a definite method cannot be established regarding aforementioned features, the V-Y flap and the Limberg flap, which are two popular methods, were compared in this study.^{6,7} There are only three studies in the English literature comparing the V-Y flap and Limberg flap.

There are three basic principles in the surgical treatment of pilonidal sinus disease. These are complete eradication of the sinuses, good closure of the remaining defective area and prevention of recurrences.^{5,8} There is no contradiction in the complete removal of the sinuses, however, the main discussion is on how to close the defect formed.^{9,10} Leaving large defects open or marsupialization are not preferred today due to long recovery time and poor cosmetic results.^{5,9,10} It is difficult to perform flap techniques with more limited dissection such as Karydakakis and Bascom without creating tension in large and complicated cases. As is well known, good wound healing can only be achieved by well-fed tissues and minimal tension. Therefore, various flap techniques have been developed and many studies have been published. While many factors such as being primary or recurrence, number of sinuses, defect size, patient preference, and surgeon experience play role in the preferred surgical technique, the most popular flap techniques for cases with large tissue loss are V-Y flap and Limberg flap techniques.

The 2-year recurrence rate of Limberg flap method, which is one of the most frequently used techniques, is reported as 0-6% in the literature.¹¹ A similar recurrence rate was found in this study (6%). Limberg technique has low recurrence rate and low hospital stay in both primary and recurrent pilonidal sinus cases. However, in some studies in which Limberg flap method has been performed, cosmetic dissatisfaction is a significant disadvantage.^{12,13} In a study in sixty-three patients, 63% of patients were not satisfied with the cosmetic result.¹³

V-Y flap technique was first used by Khatri et al.,⁶ in 1994 in five patients and it is preferred in recurrent cases. The researchers argued that V-Y flap technique provided a better wound healing due to lesser tension, lesser postoperative seroma and dead space rates as a result of limited dissection and mobilization compared to Limberg flap.^{14,15} The

disadvantage of V-Y flap method reported in the literature seems to be longer operative time and longer hospital stay.^{16,17} Operative times were not compared in our study. However, the length of hospital stay of patients in the V-Y group was similar to that of the Limberg group with a mean of 2 days and was significantly lower than the literature. Similarly, time to return to work of the patients who underwent V-Y flap was similar to the Limberg group and was shorter than the other studies in the literature.

The reported rates for V-Y flap technique was 0-10.2% for wound site infection, 0-4.6% for seroma and 0-10.2% for wound dehiscence.^{16,17,18} The rates we obtained in our study were slightly higher compared to the literature. The reason for this was concluded to be lower sociocultural levels and self-care of our patients and that they did not comply with the post-discharge recommendations.

A controversial subject in the literature is the use of drains. There are publications regarding them as beneficial and there are also studies claiming the opposite. Our general experience in pilonidal sinus surgery is the use of drains. In this study, seroma development is significantly less in all of our patients due to the use of drains. Therefore, in obese patients and those who have large tissue defects, we recommend that the drain is placed and is not removed until the amount of discharge is reduced.

Recurrence rate is one of the most important criteria in the success of surgical procedures. Both techniques have significantly less recurrence rates compared to other methods. The recurrence rate for V-Y flap and Limberg flap were 0-11% and 0-6%, respectively, in the literature.^{11,17,18} In our study, recurrence was detected in only two patients (4%) in the V-Y flap group and three patients in the Limberg group (6%) in a 2-year period. Although recurrence rate following Limberg procedure is close to the literature, our recurrence rate in the V-Y group is lower.

Conclusion

In our study, the superiority of V-Y flap technique to Limberg was observed in terms of cosmetic satisfaction and low recurrence rate. Regarding wound dehiscence, V-Y flap results seem to be better with a statistically significant difference. According to the results of this study, V-Y flap technique should be considered as a good choice for the closure of large tissue defects in the surgical treatment of pilonidal sinus disease.

Ethics

Ethics Committee Approval: The study was approved by the Yıldırım Beyazıt University Faculty of Medicine Ethics Committee (approval number: 20.01.2016/09).

Informed Consent: Written informed consent was obtained from all patients preoperatively.

Peer-review: Externally peer-reviewed.

Authorship Contributions

Surgical and Medical Practices: S.A., R.S., Concept: R.H.G., Design: R.H.G., R.S., Data Collection or Processing: R.S., S.A., Analysis or Interpretation: M.Ö.Y., Literature Search: R.H.G., R.S., Writing: R.S.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study received no financial support.

References

1. Sondenaa K, Andersen E, Nesvik I, Soreide JA. Patient characteristics and symptoms in chronic pilonidal sinus disease. *Int J Colorectal Dis* 1995;10:39-42.
2. Kallis MP, Maloney C, Lipskar AM. Management of pilonidal disease. *Curr Opin Pediatr* 2018;30:411-416.
3. Johnson EK. Expert Commentary on Pilonidal Disease: Management and Definitive Treatment. *Dis Colon Rectum* 2018;61:777-779.
4. Humphries AE, Duncan JE. Evaluation and management of pilonidal disease. *Surg Clin North Am* 2010;90:113-124.
5. De Parades V, Bouchard D, Janier M, Berger A. Pilonidal sinus disease. *J Visc Surg* 2013;150:237-247.
6. Khatri VP, Espinosa MH, Amin AK. Management of recurrent pilonidal sinus by simple V-Y fasciocutaneous flap. *Dis Colon Rectum* 1994;37:1232-1235.
7. Ersoy E, Devay AO, Aktimur R, Doganay B, Ozdogan M, Gündoğdu RH. Comparison of the short-term results after Limberg and Karydakias procedures for pilonidal disease: randomized prospective analysis of 100 patients. *Colorectal Dis* 2009;11:705-710.
8. Shabbir J, Chaudhary N.N, Britton D.C. Management of sacrococcygeal pilonidal sinus disease: a snapshot of current practice. *Int J Colorectal Dis* 2011;26:1619-1620.
9. Segre D, Pozzo M, Perinotti R, Roche B. Italian Society of Colorectal Surgery. The treatment of pilonidal disease: guidelines of the Italian Society of Colorectal Surgery (SICCR). *Tech Coloproctol* 2015;19:607-613.
10. Iesalnieks I, Ommer A, Petersen S, Doll D, Herold A. German national guideline on the management of pilonidal disease. *Langenbecks Arch Surg* 2016;401:599-609.
11. Stauffer VK, Luedi MM, Kauf P, Schmid M, Diekmann M, Wieferrich K, et al. Common surgical procedures in pilonidal sinus disease: A meta-analysis, merged data analysis, and comprehensive study on recurrence. *Sci Rep* 2018;15:8:3058.
12. Bozkurt MK, Tezel E. Management of pilonidal sinus with Limberg flap. *Dis Colon Rectum* 1998;41:775-777.
13. Eryilmaz R, Sahin M, Alimoglu O, Dasran F. Surgical treatment of sacrococcygeal pilonidal sinus with the Limberg transposition flap. *Surgery* 2003;134:745-749.
14. Schoeller T, Wechselberger G, Otto A, Papp C. Definite surgical treatment of complicated recurrent pilonidal disease with a modified fasciocutaneous V-Y advancement flap. *Surgery* 1997;121:258-263.

15. Sahasrabudhe P, Panse N, Waghmare C, Waykole P. V-Y advancement flap technique in resurfacing postexcisional defect in cases with pilonidal sinus disease-study of 25 cases. *Indian J Surg* 2012;74:364-370.
16. Berkem H, Topaloglu S, Ozel H, Avsar FM, Yildiz Y, Yüksel BC, et al. V-Y advancement flap closures for complicated pilonidal sinus disease. *Int J Colorectal Dis* 2005;20:343-348.
17. Öz B, Akcan A, Emek E, Akyüz M, Sözüer E, Akyıldız H, et al. A comparison of surgical outcome of fasciocutaneous V-Y advancement flap and Limberg transposition flap for recurrent sacrococcygeal pilonidal sinus disease. *Asian J Surg* 2017;40:197-202.
18. Altıntoprak F, Dikicier E, Arslan Y, Ozkececi T, Akbulut G, Dilek ON. Comparison of the Limberg flap with the V-Y flap technique in the treatment of pilonidal disease. *J Korean Surg Soc* 2013;85:63-67.



Evaluation of Etiological Risk Factors in the Development of Adult Chronic Pilonidal Disease

Erişkin Kronik Pilonidal Sinüs Hastalığının Oluşumunda Etiyolojik Risk Faktörlerinin Değerlendirilmesi

© Adnan Kuvvetli, © Süleyman Çetinkunar, © Alper Parlakgümüş

University of Health Sciences, Adana City Training and Research Hospital, Clinic of General Surgery, Adana, Turkey

ABSTRACT

Aim: Pilonidal disease has a high rate of recurrence after surgical interventions and measures that can be taken to prevent the recurrence are gaining importance. This study was planned to review the possible risk factors and to provide guidance for the reduction of pilonidal disease.

Method: Patients diagnosed with pilonidal disease between January 2014 and January 2017 were evaluated. Age, gender, weight, height, body mass index (BMI), occupation and sitting or standing position during working, family history of pilonidal sinus, skin color and hair color were analyzed retrospectively. The BMI of the patients was categorized as ≤ 20.0 kg/m², 20.1-25.0 kg/m², 25.1-30.0 kg/m², 30.1-35.0 kg/m² and 35.1-40.0 kg/m².

Results: The total number of patients with pilonidal disease was 217 with a mean age 26.0. The disease had two age peaks at 19 and 24-27 years. Pilonidal disease was statistically significantly more commonly observed in patients with BMI of 25.1-30.0 kg/m² ($p < 0.001$). The disease was significantly higher in brunette patients compared to auburn and blonde patients ($p < 0.001$). The disease was observed in 49.8% patients with seated positional occupations ($p < 0.001$).

Conclusion: Age, skin color and hair color are unmodifiable risk factors, but prevention of obesity, reducing sitting time and changing routines while fulfilling occupations seem preventive methods.

Keywords: Etiology, pilonidal sinus, prevention

ÖZ

Amaç: Pilonidal sinüs hastalığı cerrahisi sonrası yüksek oranda oluşan nüksler nedeniyle hastalık oluşmaması için alınabilecek önlemler önem kazanmaktadır. Bu çalışma olası risk faktörlerinin gözden geçirilip pilonidal hastalığın önlem alınarak azaltılmasında ışık tutması amacıyla planlanmıştır.

Yöntem: Ocak 2014 ve Ocak 2017 arası pilonidal sinüs tanısı almış hastalar değerlendirildi. Yaş, cinsiyet, ağırlık, boy, vücut kitle indeksi (VKİ), iş ve oturur veya ayakta icra edilişi, ailede pilonidal sinüs varlığı, deri ve saç rengi, retrospektif olarak analiz edildi. Hastaların VKİ'si $\leq 20,0$ kg/m², 20,1-25,0 kg/m², 25,1-30,0 kg/m², 30,1-35,0 kg/m² and 35,1-40,0 kg/m² olarak kategorize edildi.

Bulgular: Pilonidal sinüs olan 217 hastanın ortalama yaşı 26,0 idi. On dokuz ve 24-27 yaş aralığı, hastalarda 2 ayrı pik yapmış olarak görüldü. Pilonidal sinüs VKİ'si 25,1-30,0 kg/m² olan hastalarda istatistiksel olarak sık görüldü ($p < 0,001$). Esmer tenli hastalarda, sarışın ve kumrallara göre istatistiksel olarak bu hastalık daha sık idi ($p < 0,001$). Oturarak icra edilen mesleklerde bu hastalığın görülme oranı %49,8 idi ($p < 0,001$).

Sonuç: Hastanın yaşı, ten ve saç rengi değiştirilebilir risk faktörü değildir ancak obezitenin engellenmesi, oturur pozisyonda yapılan işlerde mümkünse sürenin kısaltılması veya icra ederken iş alışkanlığının değiştirilmesi preventif olabilir görülmektedir.

Anahtar Kelimeler: Etiyoloji, pilonidal sinüs, önlem



Address for Correspondence/Yazışma Adresi: Alper Parlakgümüş MD,
University of Health Sciences, Adana City Training and Research Hospital, Clinic of General Surgery, Adana, Turkey
Phone: +90 532 718 29 66 E-mail: aparlakgumus@yahoo.com ORCID ID: orcid.org/0000-0002-0725-6629
Received/Geliş Tarihi: 23.11.2018 Accepted/Kabul Tarihi: 25.01.2019

Introduction

Pilonidal sinus (PS) is a common surgical disease with a midline or lateral hair-containing sinus in the natal cleft.¹ Although the etiology is not known exactly, risk factors include increased hair amount in the region, male gender, obesity, prolonged sitting, family history, trauma, irritation of the coccyx, and inattention to personal hygiene.² The incidence shows variability due to geographical- and population-based conditions.^{3,4} Because of the high rates of recurrence after surgical interventions, prevention is preferred to therapy. Therefore, here we performed this study to determine the factors associated with an increased risk of PS that could be avoided, and to provide guidance for the reduction of PS.

Materials and Methods

The project was approved by the local ethics committee (project no: 325). Patients diagnosed with PS between January 2014 and January 2017 were evaluated. Age, gender, weight, height, body mass index (BMI), occupation and sitting or standing position during working, family history of PS, skin color and hair color were analyzed retrospectively. The BMI of the patients was categorized as ≤ 20.0 kg/m², 20.1-25.0 kg/m², 25.1-30.0 kg/m², 30.1-35.0 kg/m² and 35.1-40.0 kg/m². The patients with missing data, acute PS and younger than 18 years old were excluded from the study.

Statistical Analysis

The data analysis was performed using SPSS for Windows, version 22. The normality of the distribution of continuous variables was determined by Kolmogorov-Smirnov test and histogram. The data were reported as mean \pm standard deviation, or median and range, where applicable. The categorical data were analyzed using chi-square test when appropriate. A p value less than 0.05 was considered statistically significant.




Figure 1. Age distribution of patients with pilonidal disease

Results

The total number of patients with PS was 217. The patients were between the ages of 18-67 (median: 26 years) (Figure 1). The male to female ratio was 5.6. Female subjects were older at the time of diagnosis. The median ages for female and male patients were 27 and 24 years, respectively, but the age distribution was not statistically significant between two genders ($p=0.398$). The disease had two age peaks at 19 and 24-27 years. (Figure 1).

The BMI distribution among patients was seen in Figure 2. Pilonidal disease was statistically significantly more commonly observed in patients with BMI of 25.1-30.0 kg/m² ($p<0.001$).




Figure 2. Body mass index distribution of patients with pilonidal disease




Figure 3. Skin and hair color distribution of patients




Figure 4. The distribution of occupations and working positions

The color of the skin and hair was also an important factor (Figure 3). The disease was significantly higher in brunette patients compared to auburn and blonde patients ($p < 0.001$). The most important variable affecting the disease was the sitting position (Figure 4). The disease was observed in 49.8% patients with seated positional occupations ($p < 0.001$). There were no correlation between family history and occurrence of the disease ($p = 0.662$).

Discussion

The etiology of PS is a problematic issue in medicine.⁵ Most of the patients (70%) reported in the literature were between 20 and 30 years of age.⁶ In our study, the mean age of patients was 26 years. More secretion in pilosebaceous glands due to sex hormones in young population may trigger the disease. In our study, the male to female ratio was 5.6 as in previous studies⁷, which means that the disease occurs more frequently in men.

High BMI is an important factor for the occurrence of PS.⁸ In this study, PS was statistically significantly more commonly observed in patients with BMI of 25.1-30.0 kg/m² as in the study of Bolandparvaz et al.,⁷ referring that overweight patients (BMI >25) were at greater risk of disease. Deep and moist natal cleft in overweight or obese patients is more susceptible to injury with hair ends due to vacuum effect of the area.⁹

In the literature, the skin and hair colors are not well-evaluated variables. In our study, we found that brunette patients had the disease more than auburn and blonde patients significantly. From the point of view, this variable should be evaluated prospectively assessed in other studies.

The relationship between occupation and PS has been notified. In one report, Kaymakcioglu et al.,⁶ emphasized the importance of sitting position while working, as in our study.

Conclusion

In conclusion, age, skin color and hair color are unmodifiable risk factors, but prevention of obesity, reducing sitting time and changing routines while fulfilling occupations seem preventive methods.

Ethics

Ethics Committee Approval: The study was approved by the Adana City Training and Research Hospital Clinical Research Ethics Committee (approval number: 325).

Informed Consent: Retrospective study.

Peer-review: Externally peer-reviewed.

Authorship Contributions

Surgical and Medical Practices: A.K., S.C., Concept: A.P., A.K., Design: A.K., Data Collection or Processing: S.C., Analysis or Interpretation: A.K., Literature Search: A.P., Writing: A.K., A.P.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study received no financial support.

References

1. Arda IS, Guney LH, Sevmis S, Hicsonmez A. High body mass index as a possible risk factor for pilonidal sinus disease in adolescents. *World J Surg* 2005;29:469-471.
2. Humphries AE, Duncan JE. Evaluation and management of pilonidal disease. *Surg Clin North Am* 2010;90:113-124.
3. Akinci OF, Bozer M, Uzunkoy A, Duzgun SA, Coskun A. Incidence and aetiological factors in pilonidal sinus among Turkish soldiers. *Eur J Surg* 1999;165:339-342.
4. Sondenaa K, Andersen E, Nesvik I, Søreide JA. Patient characteristics and symptoms in chronic pilonidal sinus disease. *Int J Colorectal Dis* 1995;10:39-42.
5. Doll D, Friederichs J, Dettmann H, Boulesteix AL, Duesel W. Time and rate of sinus formation in pilonidal sinus disease. *Int J Colorectal Dis* 2008; 23:359-364.
6. Kaymakcioglu N, Yagci G, Simsek A, Unlu A, Tekin DF, Cetiner S, et al. Treatment of pilonidal sinus by phenol application and factors affecting the recurrence. *Tech Coloproctol* 2005;9:21-24.
7. Bolandparvaz S, Moghadam Dizaj P, Salahi R, Paydar S, Banzadeh M, Abbasi HR, et al. Evaluation of the risk factors of pilonidal sinus: a single center experience. *Turk J Gastroenterol* 2012;23:535-537.
8. Harlak A, Mentis O, Kilic S, Coskun K, Duman K, Yilmaz F. Sacrococcygeal pilonidal disease: analysis of previously proposed risk factors. *Clinics (Sao Paulo)* 2010;65:125-131.
9. Allen-Mersh TG. Pilonidal sinus: finding the right track for treatment. *Br J Surg* 1990;77:123-132.



Birth Trauma: Sphincter Injuries

Doğum Travması: Sfinkter Yaralanması

© Murat Çakır, © Mehmet Aykut Yıldırım

Necmettin Erbakan University, Meram Faculty of Medicine, Department of General Surgery, Konya, Turkey

ABSTRACT

Aim: Fecal incontinence is a condition that has social and economic effects and that disrupts the quality of life. A complete definition of fecal incontinence is difficult. However, it can be defined as the failure to control the anal discharge of intestinal content at an appropriate time and place. The diagnosis of fecal incontinence is challenging and complex. One of the most important causes of fecal incontinence is trauma in the perianal region. The most important cause of such trauma is birth trauma. We examined our cases who had sphincter repair due to acute injury in the perianal region during delivery.

Method: The data of 10 female patients who were admitted to Necmettin Erbakan University, Meram Faculty of Medicine, Department of General Surgery with perianal injuries between 2010 and 2015 were analyzed retrospectively. Data regarding age, severity of injury, first intervention time, repair type, wound problems and post-operative complications were investigated. Long-term results were analyzed.

Results: It was observed that 10 female patients underwent emergency surgery due to acute perianal injury. Ten patients were included in the study and their files were analyzed retrospectively. The mean age of the patients was 24 (range: 19-36) years. Four patients had type 4 injury, two had type 3a injury, one had type 3b injury, and three had type 3c injury.

Conclusion: We believe that it is important to perform surgical treatment before tissue edema develops.

Keywords: Childbirth, incontinence, trauma

ÖZ

Amaç: Fekal inkontinans sosyal ve ekonomik etkileri olan ve yaşam kalitesini bozan bir durumdur. Fekal inkontinansın tam bir tanımını yapmak zordur. Ancak barsak içeriğinin uygun zaman ve yerde, anal yoldan çıkışının kontrol edilememesi şeklinde tanımlanabilir. Fekal inkontinans tanı zor ve karmaşıktır. Fekal inkontinansın en önemli nedenlerinden biri perianal bölgede oluşan travmadır. Bu travmanın en önemli nedeni doğum travmasıdır. Doğum esnasında perianal bölgede oluşan akut yaralanma nedeniyle sfinkter tamiri yapılan olgularımızı inceledik.

Yöntem: Necmettin Erbakan Üniversitesi Meram Tıp Fakültesi Genel Cerrahi Kliniği 2010-2015 yılları arasında başvuran perianal bölge yaralanması olan 10 kadın hastanın verileri retrospektif olarak incelendi. Hastaların yaş, yaralanma derecesi, ilk müdahale zamanı, onarım şekli, yara problemleri ve postoperatif komplikasyonları değerlendirildi. Uzun dönem sonuçları incelendi.

Bulgular: On kadın hastanın akut perianal bölge yaralanması nedeniyle acil cerrahi uygulandığı tespit edildi. Çalışmaya 10 hasta alındı ve dosyaları retrospektif olarak incelendi. Kadın hastalarının yaş ortalaması 24 (19-36) idi. Kadın hastaların 4'ünde tip 4 yaralanma, 2'sinde tip 3a, 1'inde tip 3b ve 3'ünde tip 3c yaralanma izlendi.

Sonuç: Cerrahi müdahalenin doku ödemi gelişmeden yapılmasının önemli olduğu kanaatindeyiz.

Anahtar Kelimeler: Doğum, inkontinans, travma

Introduction

Fecal incontinence is a condition that has social and economic effects and disrupts quality of life.¹ It is hard to offer an exact definition of fecal incontinence. However, it

can be defined as the failure to control the anal discharge of intestinal content at an appropriate time and place.² The prevalence, diagnosis, and treatment methods of fecal incontinence could not be clearly identified. Many methods used in diagnosis and treatment have evolved over time.³ The



Address for Correspondence/Yazışma Adresi: Mehmet Aykut Yıldırım MD,
Necmettin Erbakan University, Meram Faculty of Medicine, Department of General Surgery, Konya, Turkey
E-mail: drmayildirim@hotmail.com ORCID ID: orcid.org/0000-0001-9580-6004
Received/Geliş Tarihi: 31.12.2018 Accepted/Kabul Tarihi: 03.02.2019

diagnosis of fecal incontinence is challenging and complex. One of the most significant causes of fecal incontinence is trauma in the perianal region. The most important cause of such trauma is birth trauma. It is difficult to evaluate patients and to plan treatment. The surgical method is determined by the severity of sphincter damage and injury to organs in the perianal region. Therefore, the aim of this study was to analyze the patients who underwent sphincter repair due to acute injuries in the perianal region.

Materials and Methods

The data of 10 female patients who were admitted to Necmettin Erbakan University, Meram Faculty of Medicine, Department of General Surgery with perianal injuries between 2010 and 2015 were analyzed retrospectively. Data regarding age, severity of injury, first intervention time, repair type, wound problems and post-operative complications were investigated. Long-term results were analyzed. Patients with acute injury requiring emergency intervention were included in the study. Patients who did not undergo emergency intervention for a reason but who underwent repair procedures in the long-term were excluded from the study.

Results

It was observed that 10 patients underwent emergency surgery due to acute perianal injury. It was determined that 10 patients experienced injury during childbirth. Ten patients were included in the study and their files were analyzed retrospectively. The mean age of the patients was 24 (range: 19-36) years. Four patients had type 4 injury, two had type 3a injury, one had type 3b injury, and 3 had type

3c injury (Figure 1). Six patients were primipares and six patients gave birth to babies weighing more than 4000 gr. Six patients had episiotomy and 3 did not receive episiotomy. Moreover, six of the patients had vacuum assisted labor. Nine of the patients had emergency intervention within the first 10 (range: 1-18) hours. One patient was initially treated at an external center and presented to our center on the 3rd day upon split formation in the injury. This patient was re-operated (Table 1).

While all patients received sphincteroplasty and vaginoplasty, five patients underwent levatorplasty. After repair, the rectovaginal septum was thicker than 2 cm in all patients. Three/0 polyglactin sutures were used for all repairs and interrupted repair was completed (Figure 2). Drains were placed in the repair area. Vaginal tampons were used and were frequently changed to reduce the risk of surgical site complications. Colorectal irrigation was performed in all patients during the surgical procedure.

Colostomy was not created for any patient, however, medical ileus was created. Parenteral feeding was continued for six days. All patients received third generation cephalosporin and metronidazole. Oral intake was enabled on the 5th day and the patients were discharged between the 7th and 10th days. Patients were asked to avoid sexual intercourse for 3 months. They were advised not to undergo vaginal delivery in their following labors. All patients received laxatives in

Table 1. Patient data

Variable	
Number	10
Age, years	24 (19-36)
Severity of injury	Type 4 (n=4)
	Type 3a (n=2)
	Type 3b (n=1)
	Type 3c (n=3)
Pre-operative time	10 (1-18) hour (n=1, 3 day)
Postoperative follow-up time	3 (1-7) years
Mean resting pressure	40 (30-70) mmHg
Mean extrusion pressure	60 (50-120) mmHg
Mean duration of extrusion	30 (20-56) seconds
Mean coughing reflex	70 (54-110) mmHg



Figure 1. A patients with type 3c injury



Figure 2. Post-repair image

order to ease stool discharge and to prevent constipation for a month. This period was extended in patients with chronic constipation.

All patients were followed-up for one month. None of the patients had wound site pathologies. Anal manometric pressure measurements were conducted at the 3rd month. The mean resting pressure was 40 (range: 30-70) mmHg, extrusion pressure was 60 (range: 50-120) mmHg, mean duration of extrusion was 30 (range: 20-56) seconds, and coughing reflex was 70 (range: 54-110) mmHg.

Three patients experienced gas incontinence at the 6th month. These patients received biofeedback for 2 months upon failure to detect complete sphincter damage by imaging techniques. Complaints of the patients improved after two months. The patients have been in the follow-up program for a mean period of 3 years and they do not have complaints of incontinence.

Discussion

The main causes of the problem in most of the patients with fecal incontinence are obstetric trauma or sphincter defects that emerge after anal surgery. There are varying degrees of injury in 53% to 79% of patients following vaginal delivery.^{4,5} Although the rate of such injuries is too high, not all injuries cause sphincter damage. The rate of primary obstetric anal sphincter injury is approximately 18% in vaginal deliveries.⁶ Eight studies were investigated in a meta-analysis and it was seen that 5.7% of women had anal sphincter injury during first labor and 6.3% during the following labors.⁷ In another study, a total of 700 patients pregnant with twins were evaluated and the rate of sphincter injury was found to be 2.8%.⁸ In a study of patients who had vaginal or C-section deliveries, the authors found that complaints related to incontinence varied between 13% and 25%.⁹ The rate of fecal incontinence in patients with postpartum sphincter injury is approximately 7.7%.¹⁰ While the rate of incontinence is 6.3% in nulliparous women, it is 8.8% in uniparous women and 8.4% in secundiparous women, compared with 11.5% in triparous women and women who had more than 3 deliveries.¹¹ Only 27% of endoanal ultrasonography results show occult anal sphincter injuries.¹² One third of the patients with detected damage also had symptoms of incontinence. Asymptomatic patients are at risk for incontinence in the coming years. All of our patients had anal sphincter damage due to birth trauma. The vast majority of our cases sustained anal sphincter injury during the first delivery.

Creation of episiotomy during labor prevents uncontrolled perianal fissures. However, the results of a meta-analysis is revealed that episiotomy increased the risk of fecal

incontinence.¹³ The most important parameters for sphincter damage at vaginal delivery are interventional delivery and birthweight of the baby.¹⁴ Tertiary fissures are frequently seen in interventional deliveries with midline episiotomy. Uncontrolled fissures are less common after medio-lateral episiotomy procedures. Although six patients had episiotomy, uncontrolled fissures could not be prevented. We believe this was due to the problematic timing of episiotomy. Although the starting point of the episio-incision was medio-lateral, it then proceeded medially. Six patients had interventional deliveries. Six of the babies had a birth weight of more than 4000 grams.

The first treatment that should be offered for any incontinence patient with anal sphincter defect is primary sphincter repair. Pre-operative intestinal cleaning should be done very carefully. Perioperative antibiotic prophylaxis is necessary. Randomized controlled studies have demonstrated that prophylaxis with second-generation cephalosporin significantly reduces the rate of infection.¹⁵ Infection rates following anal sphincter repair due to vaginal birth injury is as high as 20%, whereas, wound healing problems reach up to 25%.¹⁶ Surgical site infection was not observed in any of our patients. We administer a combination of third generation cephalosporin and metronidazole prophylaxis, and we continue treatment for 3 days. We pay special attention to keep the vagina dry. We believe that the reason why we do not have complications such as wound site infection and wound splitting is the attention we pay.

The vaginal wall on the front and anal canal mucosa on the back should be repaired in internal and external sphincter procedures. Sphincter repairs can be done in end-to-end or overlapping forms. The difference between end-to-end and overlapping suturing has not been shown in the literature within one-year of follow-ups.¹⁷ We do end-to-end repair in cases with early surgery following injury. We prefer the overlapping method more for chronic cases. We repair all layers separately with interrupted sutures.

There is no sufficient data on whether a patient who had vaginal injury following vaginal delivery should have her next deliveries by vaginal or C-section delivery. However, some studies have reported that the possibility of a new anal sphincter injury following vaginal delivery is 3%.¹⁸ Nevertheless, we do not recommend vaginal delivery to our patients. These patients have lower anal manometric pressure than normal patients (although the measurements do not have standard values). Micro or macro sphincter injuries happen in all deliveries. We support the idea that patients should not be exposed to the risk of having possible injuries.

Incontinence can be seen in women with sphincter injuries during delivery after repair. The most important cause of this

is residual anterior sphincter defects.³ Adequate continence might not be achieved in the long term by the primary repair of third and fourth degree sphincter fissures that form during delivery. The incidence of anal incontinence was reported as 15-61%, although primary repair is done during delivery. The endoanal ultrasonography results presented in some studies have demonstrated that sphincter injuries remain up to 91% in patients with fissures repaired by interrupted side-to-side sutures.^{3,19} Three-year follow-up results of our patients have not shown any significant continence problems. Biofeedback therapy has been shown to be helpful for patients suffering from gas incontinence and urogenesis problems.

Conclusion

Physicians should note that sphincter injuries might happen after every delivery. Patients should be informed about the fact that they might experience incontinence problems during the follow-ups. We believe that it is significant to perform surgical intervention before tissue edema develops.

Ethics

Ethics Committee Approval: The study was approved by the Necmettin Erbakan University Meram Faculty of Medicine Ethics Committee (approval number: 2017/292).

Informed Consent: Informed consent was obtained from all individual participants included in the study.

Peer-review: Externally peer-reviewed.

Authorship Contributions

Surgical and Medical Practices: M.Ç., M.A.Y., Concept: M.Ç., Design: M.Ç., Data Collection or Processing: M.Ç., Analysis or Interpretation: M.Ç., M.A.Y., Literature Search: M.Ç., M.A.Y., Writing: M.Ç., M.A.Y.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study received no financial support.

References

- Landefeld CS, Bowers BJ, Feld AD, Hartmann KE, Hoffman E, Ingber MJ, et al. National Institutes of Health state-of-the-science conference statement: prevention of fecal and urinary incontinence in adults. *Ann Intern Med* 2008;148:449.
- Wald A, Bharucha AE, Cosman BC, Whitehead WE. ACG clinical guideline: management of benign anorectal disorders. *Am J Gastroenterol* 2014;109:1141.
- Bulut MT, Fekal inkontinans. In: Menteş B, Bulut MT, Alabaz Ö, Leventoglu S. *Anorektal Bölgenin Selim Hastalıkları*. 1. Baskı, Ankara, 2011:187-216.
- Smith LA, Price N, Simonite V, Burns EE. Incidence of and risk factors for perineal trauma: a prospective observational study. *BMC Pregnancy Childbirth* 2013;7;13:59. doi: 10.1186/1471-2393-13-59.
- Rogers RG, Leeman LM, Borders N, Qualls C, Fullilove AM, Teaf D, et al. Contribution of these condstage of labourto pelvic floor dysfunction: a prospective cohort comparison of nulliparous women. *BJOG* 2014;121:1145-1153.
- Lowder JL, Burrows LJ, Krohn MA, Weber AM. Risk factors for primary and subsequent anal sphincter lacerations: a comparison of cohorts by parity and prior mode of delivery. *Am J Obstet Gynecol* 2007;196:344.e1
- Jha S, Parker V. Risk factors for recurrent obstetric anal sphincter injury (rOASI): a systematic review and meta-analysis. *Int Urogynecol J* 2016;27:849-57. Epub 2015 Dec 16.
- Rosen H, Barrett J, Okby R, Nevo O, Melamed N. Risk factors for obstetric anal sphincter injuries in twin deliveries: a retrospective review. *IntUrogynecol J* 2016;27:757-62. Epub 2015 Sep 30.
- Guise JM, Morris C, Osterweil P, Li H, Rosenberg D, Greenlick M. Incidence of fecal incontinence after childbirth. *Obstet Gynecol* 2007;109:281.
- Melville JL, Fan MY, Newton K, Fenner D. Fecal incontinence in US women: a population-basedstudy. *Am J Obstet Gynecol* 2005;193:2071.
- Nygaard I, Barber MD, Burgio KL, Kenton K, Meikle S, Schaffer J, et al. Pelvic Floor Disorders Network. Prevalence of symptomatic pelvic floor disorders in US women. *JAMA* 2008;300:1311.
- Oberwalder M, Connor J, Wexner SD. Meta-analysis to determine the incidence of obstetric anal sphincter damage. *Br J Surg* 2003;90:1333-1337.
- LaCross A, Groff M, Smaldone A. Obstetric anal sphincter injury and anal incontinence following vaginal birth: a systematic review and meta-analysis. *J Midwifery Womens Health* 2015;60:37-47.
- Pergialiotis V, Vlachos D, Protopapas A, Pappa K, Vlachos G. Risk factorsfor severe perineal lacerations during childbirth. *Int J Gynaecol Obstet* 2014;125:6-14.
- Duggal N, Mercado C, Daniels K, Bujor A, Caughey AB, El-Sayed YY. Antibiotic prophylaxis for prevention of postpartum perineal wound complications: a randomized controlled trial. *Obstet Gynecol* 2008;111:1268-1273.
- Lewicky-Gaupp C, Leader-Cramer A, Johnson LL, Kenton K, Gossett DR. Wound complications after obstetric anal sphincter injuries. *Obstet Gynecol* 2015;125:1088-1093.
- Fernando RJ, Sultan AH, Kettle C, Thakar R. Methods of repair for obstetric anal sphincter injury. *Cochrane Database of Systematic Reviews* 2013;12: CD002866. DOI: 10.1002/14651858.CD002866.
- Peirce C, Murphy C, Fitzpatrick M, Cassidy M, Daly L, O'Connell PR, et al. Randomised controlled trial comparing early home biofeedback physiotherapy with pelvic floor exercises for thetreatment of third-degree tears (EBAPT Trial). *BJOG* 2013;120:1240-1247.
- Practice Bulletin No. 165: Prevention and Management of Obstetric Lacerations at Vaginal Delivery. American College of Obstetricians and Gynecologists' Committee on Practice Bulletins-Obstetrics. *Obstet Gynecol* 2016;128:e1-e15.



An Acute Abdomen Dilemma: Epiploic Appendagitis

Bir Akut Karın Dilemması: Epiploik Appendajit

© Hakan Özdemir¹, © Oğuzhan Sunamak¹, © Zehra Ünal Özdemir¹, © Ferdi Cambaztepe²

¹İstanbul Haydarpaşa Numune Training and Research Hospital, Clinic of General Surgery, İstanbul, Turkey

²Kaçkar State Hospital, Clinic of General Surgery, Rize, Turkey

ABSTRACT

Aim: Appendagitis is a clinical condition caused by ischemia, torsion or inflammation of epiploic appendices located on serosal surface of the colon. Antibiotics and analgesics are generally sufficient in treatment. Rarely, excision might be needed. It might be confused with acute appendicitis and diverticulitis, depending on its localization.

Method: The data of 12 patients with acute abdomen, who were diagnosed to have epiploic appendagitis and responded to medical treatment completely, were analyzed retrospectively.

Results: There were seven female and five male patients with a mean age of 33 (range: 21-48) years. The mean body mass index was 25.5 (range: 19-34). There was no abdominal surgery. The mean length of hospital stay, leukocyte count and C-reactive protein (CRP) were 2.08 days, $10.41 \times 10^3/\mu\text{L}$ and 2.3 mg/dL, respectively. Sixty-six point six percent (n=8) of the epiploic appendagitis was localized in the right colon and 33.3% (n=4) in the left colon. A positive correlation was found between the diameter of appendagitis and leukocyte count and CRP level ($p>0.05$). There was no correlation between appendagitis diameter and vomiting ($p>0.05$).

Conclusion: Appendagitis should be kept in mind in patients presenting with sudden onset, sharp and constant pain. Informing radiologist about this possibility may help to confirm the diagnosis.

Keywords: Conservative treatment, epiploic appendagitis, radiology

ÖZ

Amaç: Apandijit, kolon serozasında bulunan appendiks epiploikaların iskemi, torsiyon veya enflamasyonuna bağlı klinik durumdur. Antibiyotik ve analjezik tedavi tedavide genellikle yeterlidir. Nadiren, eksizyon gerekebilir. Yerleşim yerine bağlı olarak, akut apandisit vedivertikülit ile karışabilir.

Yöntem: Akut karın ile başvuran ve epiploik apandijit tanısı alarak medikal tedavi ile tamamen iyileşen 12 hastanın verileri retrospektif olarak incelendi.

Bulgular: Yaş ortalaması 33 (21-48) olan yedi kadın, beş erkek hasta vardı. Ortalama vücut kitle endeksi 25,5 (19-34) idi. Hiçbir hasta ameliyat edilmedi. Ortalama hastanede kalış süresi, lökosit sayısı ve C-reaktif protein (CRP) değerleri sırasıyla, 2,08 gün, $10,41 \times 10^3/\mu\text{L}$ ve 2,3 mg/dL idi. Epiploik apandijitlerin %66,6'sı (n=8) sağ kolon ve %33,3'ü (n=4) sol kolon yerleşimliydi. Apandajit çapı, lökosit sayısı ve CRP düzeyi arasında pozitif bir ilişki saptandı ($p<0,005$). Apandajit çapı ile kusma arasında bir ilişki saptanmadı ($p>0,005$).

Sonuç: Epiploik apandajit, ani başlayan, keskin ve aralıksız ağrısı olan hastalarda akla gelmelidir. Bu olasılık açısından radyoloğun bilgilendirilmesi tanıyı kesinleştirmede yardımcı olabilir.

Anahtar Kelimeler: Konservatif tedavi, epiploik apandajit, radyoloji

Introduction

Acute primary epiploic appendagitis develops due to ischemia, torsion or inflammation of epiploic appendices located on serosal surface of the colon. Clinically, it generally presents with a sudden-onset, sharp and constant pain. It mimics acute appendicitis and diverticulitis depending on its localization, either in the right or left colon. Radiological

findings may be confused with omental infarct, panniculitis or fat-containing tumors.¹ Radiologically, ultrasonography and computed tomography are sufficient to diagnose.² If it cannot be diagnosed radiologically, patients may undergo surgery due to acute abdomen. Appendagitis is frequently a self-limiting, benign clinical condition. It is mostly seen between 2nd and 5th decades, and is more frequent in



Address for Correspondence/Yazışma Adresi: Hakan Özdemir MD,
İstanbul Haydarpaşa Numune Training and Research Hospital, Clinic of General Surgery, İstanbul, Turkey
Phone: +90 532 468 53 80 E-mail: hakanzdmr@yahoo.com ORCID ID: orcid.org/0000-0002-6304-3118
Received/Geliş Tarihi: 30.11.2018 Accepted/Kabul Tarihi: 20.02.2019

women and obese people. A careful history taking, physical examination and radiological imaging can reveal accurate diagnosis. Antibiotics and analgesics are generally sufficient for treatment.

Materials and Methods

Twelve patients who were admitted to emergency department with acute abdomen and diagnosed with epiploic appendagitis between January 2014 and November 2017 were retrospectively analyzed. Physical examination, laboratory and imaging results were recorded. Local ethical committee approved the study and informed consent was obtained from the patients.

Statistical Analysis

SPSS 17 program was used for statistical analysis. Pearson correlation test was used to compare parameters. $P < 0.05$ was considered statistically significant.

Results

There were seven female and five male patients with a mean age of 33 (range: 21-48) years. The mean body mass index (BMI) was 25.5 (range: 19-34). There was no abdominal surgery. The mean length of hospital stay, leukocyte count and C-reactive protein (CRP) were 2.08 days; $10.41 \times 10^3/\mu\text{L}$ and 2.3mg/dL, respectively. The diagnosis was made by computerized tomography (CT) scan in all patients (Figure 1 and 2). Most (66.6%) of the epiploic appendagitis was localized in the right colon (Table 1). A positive correlation was found between the diameter of appendagitis and leukocyte count and CRP level ($p < 0.05$). There was no relationship between appendagitis diameter and vomiting. The symptoms of the patients regressed with medical treatment. There was no complication or need for any surgical intervention.

Discussion

Epiploic appendices are lipomatous structures having a small arterial supply and venous return to colonic straight vessels. Matos and Costa³ was the first to describe epiploic appendagitis in 1956 in the differential diagnosis of sudden-onset, right or left lower quadrant abdominal pain.

The patients describe a sharp, sudden onset, constant abdominal pain without nausea and vomiting or fever. The pain was reported to be more frequent in the left lower quadrant in the literature.^{3,4,5} However, it was more frequent in the right lower quadrant in our study. Routine laboratory tests are within normal limits, except for mild leukocyte count and CRP increase.⁶ Before the widespread availability of CT, appendagitis could only be diagnosed during




Figure 1. Epiploic appendagitis on computerized tomography




Figure 2. Epiploic appendagitis on computerized tomography

Table 1. Demographic characteristics of patients

Patient	Gender	Age (years)	Nausea/vomiting	Pain localization	Leukocyte count ($\times 10^3/\mu\text{L}$)	CRP	Appendagitis diameter	BMI	Hospital stay (days)
1	Female	41	0	Right	7.06	0.5	1.6	20	1
2	Male	28	0	Right	10.10	2.1	1.7	27	2
3	Male	21	1	Right	9.10	2.9	2.1	23	3
4	Female	25	1	Left	10.90	3.3	2.4	32	3
5	Female	38	0	Right	13.00	1.7	1.5	29	2
6	Male	24	0	Left	8.10	2.2	1.1	19	1
7	Female	37	0	Right	12.30	4.1	2.7	33	3
8	Female	45	0	Right	11.40	1.2	1.8	25	2
9	Female	22	0	Right	9.40	1.6	2.3	27	3
10	Male	29	1	Left	11.10	1.4	1.9	34	1
11	Male	48	1	Left	13.80	5.1	3.7	23	3
12	Female	40	0	Right	8.70	2.3	1.2	31	1

CRP: C-reactive protein, BMI: Body mass index

surgical exploration, as its symptoms are non-specific. A study reported 2.5% preoperative correct diagnosis rate for appendagitis, which is very low.¹ Suspicion of appendagitis and informing the radiologist will be useful in the correct diagnosis during preoperative evaluation. Cases with non-radiological diagnosis or non-regressive symptoms despite medical therapy can undergo surgery with a prediagnosis of acute abdomen.^{7,8}

It appears as an oval, non-compressible and hypoechoic lesion in ultrasound imaging.^{9,10} There is severe tenderness on the lesion location. CT reveals a lesion of lipid density along with inflammatory findings in the neighborhood of colon.¹¹ Parietal peritoneum may appear thickened, if inflammation expands. Generally, the colonic wall appears to have normal thickness in CT, although it may appear thicker. The radiological findings of appendagitis resolve completely in 6 weeks.¹²

Epiploic appendagitis is a benign, self-limiting clinical condition with a size of 0.5 to 5 cm, and it is generally seen in young and middle-aged patients and obese people.² The high mean BMI and mean age in our study are consistent with the literature. Heavy physical activity may increase the risk of developing appendagitis. Male or female dominance is not certain in the literature.^{1,3} It may be confused with acute appendicitis, acute cholecystitis or diverticulitis, depending on localization. The patients with a definite diagnosis based on imaging should be initially treated medically. Diagnostic laparoscopy can be used in cases with unexplained clinical or non-regressive symptoms. In addition, laparoscopic excision of appendagitis can eliminate the pathology.¹³

Epiploic appendagitis is one of the blamed factors in etiology of intraperitoneal loose body, along with omental torsion and peritoneal debris.¹⁴ These intraperitoneal structures can mimic pseudotumor in imaging. Following acute period, appendagitis may show calcification and may be confused with carcinomatosis radiologically.⁷ In such situations, if available, comparison with previous radiological imaging is helpful in diagnosis; otherwise, biopsy may be needed. While necrotized epiploic appendix tissue generally shows eggshell calcification, metastatic calcifications frequently show nodular formation.¹⁵ However, in some cases, this discrimination is not clear. Thus, previous radiological imaging showing appendagitis is very important to differentiate this pathology from malignancy when suspected.

Conclusion

In conclusion, appendagitis is a clinical condition that is diagnosed by radiological imaging and that is generally conservatively treated. Ultrasound and CT scan are sufficient to make a diagnosis because of typical radiological findings. This typical appearance is important to differentiate it from acute appendicitis, omental infarction, diverticulitis and cholecystitis. Appendagitis should be kept in mind in patients with sudden onset or severe abdominal pain and mild increase in leukocyte or CRP counts. Informing

the radiologist about suspected appendagitis may facilitate diagnosis.

Ethics

Ethics Committee Approval: The study was approved by the İstanbul Haydarpaşa Numune Training and Research Hospital Ethics Committee (approval number: (07/06/2018-62977267-000-8436)).

Informed Consent: Informed consent of all patients were taken.

Peer-reviewed: Externally peer-reviewed.

Authorship Contributions

Surgical and Medical Practices: H.Ö., Z.Ü.Ö., O.S., Concept: H.Ö., Z.Ü.Ö., Design: H.Ö., Data Collection or Processing: H.Ö., Z.Ü.Ö., F.C., Analysis or Interpretation: H.Ö., O.S., Literature Search: H.Ö., F.C., Writing: H.Ö., Z.Ü.Ö., O.S.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study received no financial support.

References

1. Singh AK, Gervais DA, Hahn PF, Sagar P, Mueller PR, Novelline RA. Acute epiploic appendagitis and its mimics. *Radiographics* 2005;25:1521-1534.
2. Nadida D, Amal A, Ines M, Makram M, Amira M, Leila BF, et al. Acute epiploic appendagitis: Radiologic and clinical features of 12 patients. *Int J Surg Case Rep* 2016;28:219-222.
3. Matos H, Costa I. Primary epiploic appendagitis. *Indian J Surg* 2015;77(Suppl 3):1395-1396.
4. Almuhanha AF, Alghamdi ZM, Alshammari E. Acute epiploic appendagitis: A Rare cause of acute abdomen and a diagnostic dilemma. *J Family Community Med* 2016;23:48-50.
5. Michael A, Zakry Y, Hanif H. Epiploic appendagitis: an uncommon surgical diagnosis. *Med J Malaysia* 2017;72:128-129.
6. Ortega-Cruz HD, Martinez-Souss J, Acosta-Pumarejo E, Toro DH. Epiploic appendagitis, an uncommon cause of abdominal pain: A case series and review of the literature. *P R Health Sci J* 2015;34:219-221.
7. Eberhardt SC, Strickland CD, Epstein KN. Radiology of epiploic appendages: acute appendagitis, post-infarcted appendages, and imaging natural history. *Abdom Radiol (NY)* 2016;41:1653-1665.
8. Schnedl WJ, Krause R, Wallner-Liebmann SJ, Tafeit E, Mangge H, Tillich M. Primary epiploic appendagitis and successful outpatient management. *Med Sci Monit* 2012;18:CS48-51.
9. Özdemir H, Özdemir ZÜ, Şenol M, Şahiner İT. A rare pathology mimicking acute appendicitis: Epiploic appendagitis. *Sakarya Med J* 4;135-138.
10. Li L, Shen Z, Xu L, Guo G, Qin Y. Recurrent abdominal pain and fever as clinical manifestations: epiploic appendagitis. *Int J Clin Exp Med* 2014;7:4523-4525.
11. Menozzi G, Maccabruni V, Zanichelli M, Massari M. Contrast-enhanced ultrasound appearance of primary epiploic appendagitis. *J Ultrasound* 2014;17:75-76.
12. Singh AK, Gervais DA, Hahn PF, Rhea J, Mueller PR. CT appearance of acute appendagitis. *AJR Am J Roentgenology* 2004;183:1303-1307.
13. Vazquez-Frias JA, Castaneda P, Valencia S, Cueto J. Laparoscopic diagnosis and treatment of an acute epiploic appendagitis with torsion and necrosis causing an acute abdomen. *JSLs* 2000;4:247-250.
14. Murat FJ, Gettman MT. Free-floating organized fat necrosis: rare presentation of pelvic mass managed with Laparoscopic techniques. *Urology* 2004;63:176-177.
15. Agarwal A, Yeh BM, Brieman RS, Quayyum A, Coakley FV. Peritoneal calcification: causes and distinguishing features on CT. *Am J Roentgenol* 2004;182:441-445.



The Effects of Glutamine and N-Acetylcysteine on Experimental Colitis Induced by 2,4,6-Trinitrobenzene Sulphonic Acid in Rats

Ratlarda 2,4,6-Trinitrobenzeno Sülfonik Asitle Oluşturulan Deneysel Kolit Üzerine Glutamine ve N-Asetilsisteinin Etkisi

İsmail Cem Eray¹, Orçun Yalav¹, Kubilay Dalcı¹, Ahmet Rencüzoğulları¹, Ahmet Gökhan Sarıtaş¹, Ömer Alabaz¹, Şule Menziletoğlu Yıldız², Gülşah Seydaoğlu³, Figen Doran⁴

¹Çukurova University Faculty of Medicine, Department of General Surgery, Adana, Turkey

²Çukurova University Faculty of Medicine, Department of Medical Biochemistry, Adana, Turkey

³Çukurova University Faculty of Medicine, Department of Biostatistics, Adana, Turkey

⁴Çukurova University Faculty of Medicine, Department of Pathology, Adana, Turkey

ABSTRACT

Aim: Various factors including free oxygen radicals have been described in the pathogenesis of inflammatory bowel disease. The aim of this study was to investigate the effects of antioxidant N-acetylcysteine (NAC) and combined glutamine (Gln) on experimental colitis.

Method: In order to achieve this, rats were randomized and administered intraperitoneal NAC or oral Gln + intraperitoneal NAC one day after experimental colitis model induced with trinitrobenzen sulphonic acid. After the treatment, the rats were sacrificed and colon damage was evaluated macroscopically and microscopically. Malonyldialdehyde level and superoxide dismutase activity were analyzed biochemically.

Results: NAC and Gln + NAC groups showed macroscopically better results than the colitis group. Histopathologically, all treated groups provided significantly lower scores than the colitis group. Similarly, all treated groups provided better biochemical results than the colitis group.

Conclusion: According to the results of the experiment, it was concluded that NAC had positive effects on experimental colitis model.

Keywords: Experimental colitis, glutamine, N-acetylcysteine

ÖZ

Amaç: Enflamatuvar barsak hastalığı patogenezinde, serbest oksijen radikallerinin de aralarında bulunduğu çeşitli etkenler tanımlanmıştır. Bu çalışmanın amacı glutamin (Gln) ile kombine edilen antioksidan N-asetilsisteinin (NAC) deneysel kolit modeli üzerindeki etkilerini incelemektir.

Yöntem: Trinitrobenzeno sülfonik asit ile deneysel kolit modeli oluşturulduktan bir gün sonra ratlar randomize edilerek intraperitoneal NAC veya oral Gln + intraperitoneal NAC uygulandı. Tedavi sonrası ratlar sakrifiye edilerek, kolon hasarı makroskobik ve mikroskobik olarak değerlendirildi. Malonil dialdehid seviyesi ve süperoksit dismutaz aktivitesi biyokimyasal olarak analiz edildi.

Bulgular: NAC veya Gln + NAC verilen gruplarda makroskobik olarak kolit grubuna göre daha iyi sonuçlar elde edildi. Histopatolojik olarak tedavi verilen tüm gruplarda kolit grubuna kıyasla anlamlı olarak daha düşük skorlar elde edildi. Biyokimyasal olarak tedavi verilen tüm gruplarda kolit grubuna kıyasla daha iyi sonuçlar elde edildi.

Sonuç: Deneyden elde edilen sonuçlara göre, NAC uygulanmasının deneysel kolit modeli üzerine olumlu etkili olduğu sonucuna varıldı.

Anahtar Kelimeler: Deneysel kolit, glutamin, N-asetilsistein



Address for Correspondence/Yazışma Adresi: İsmail Cem Eray MD,

Çukurova University Faculty of Medicine, Department of General Surgery, Adana, Turkey

Phone: +90 532 553 55 03 E-mail: cem_eray@yahoo.com ORCID ID: orcid.org/0000-0002-1560-7740

Received/Geliş Tarihi: 07.01.2019 Accepted/Kabul Tarihi: 28.02.2019

Introduction

Inflammatory bowel disease (IBD) is a systemic disease that involves the gastrointestinal system and affects the quality of life. Although there are a large number of agents currently blamed for the etiology of the disease, it is more widely accepted that the immunoregulation deficiencies of the gastrointestinal tract and oxidative stress are effective and more attention is paid to free oxygen radicals.¹ In some studies, it was found that the colon biopsies of the patients with IBD had increased free oxygen radicals, and decreased glutathione (GSH) levels and antioxidant enzyme activity in patients with ulcerative colitis.² On the other hand, another study revealed that abnormally low levels of cysteine were measured IBD-related inflammation.³

New treatment methods are being searched due to developed tolerance and side effects of the drugs. Trinitrobenzene sulfonic acid (TNBS) is a substance used to create experimental colitis model.^{4,5,6,7,8} In this model, many histopathological and clinical features of Crohn's disease can be observed as in humans and the period of chronic inflammatory response can be extended up to 8 weeks. Ethanol is given to eliminate the barrier of mucous membranes without making any morphological changes in the colon and thus helps the TNBS to penetrate into the intestinal wall.⁹

The purpose of this study was to have a better understanding the role of oxidative stress in the experimental colitis model and examine the effects of NAC and glutamine (Gln) in order to develop other treatment methods. Tissue damage was evaluated macroscopically and microscopically. Superoxide dismutase (SOD) activity and malonildialdehyde (MDA) levels were measured for the evaluation of antioxidant capacity.

Materials and Methods

The protocol was approved by the Committee of Animal Care and Use, Faculty of Medicine, Çukurova University. In this study, 50 male Wistar albino rats weighing between 220 and 320 gr and with a mean age of 4 months were used. Light-dark cycle (12 h-12 h) was used during the experiment, and rats were placed in individual cages and were kept at room temperature at about 21°C. During the study, the rats were fed with distilled tap water and standard pellet feed, and the groups receiving glutamine were fed with 1g/kg of Gln mixed with distilled tap water and standard pellet. Anesthesia in rats was achieved by injecting 50 mg/kg dose of intramuscular ketamine hydrochloride (Ketalar, Parke Davis and Eczacıbaşı, İstanbul) +5 mg/kg of xylazine hydrochloride.

The Induction of Colitis

To create an experimental colitis, a mixture of 30 mg (80 mg/kg) TNBS (92823, picrylsulfonic acid solution 10 mL, purum, Switzerland) +30% ethanol was used. According to this method, 8-F polyurethane cannula was placed 8 cm rectally in anesthetized rats. Colitis was induced by administering 80 mg/kg TNBS +0.3 mL of 30% ethanol slowly through this cannula. The TNBS-E that remained in the cannula was introduced into the rectum by 1 cc air. Then, rats were held upsidedown for 30 seconds in order to prevent the reflux of the substance given and they were kept in the Trendelenburg position until they awakened from anesthesia.

Experimental Groups and Treatments (Table 1)

Group I (control group; n=10): Enema was applied with normal saline to the normal rats on the first and second days. Intraperitoneal sulforaphane (SF) was administered between the first and fourth days.

Group II (ethanol group; n=10): To monitor the possible effect of ethanol on the intestinal mucosa, 0.3 mL 30% ethanol was given through the anal canal in the supin position to a depth of about 8 cm through a thin nutrition catheter. The following day, intraperitoneal SF was given between the first and fourth days.

Group III (colitis group; n=10): A mixture of 80 mg/kg TNBS and 1 mL/kg 30% ethanol was given to a depth of about 8 cm through a thin nutrition catheter. The following day, intraperitoneal SF was given between the first and fourth days after the operation.

Group IV (NAC; n=10): A mixture of 80 mg/kg TNBS and 1 mL/kg 30% ethanol was given to a depth of about 8 cm through a thin nutrition catheter. The following day, 300 mg/kg intraperitoneal NAC injection was performed for 4 days.

Group V (Gln-NAC; n=10): Two weeks before the induction of colitis, 10 g/L Gln was mixed with drinking water. Drinking water was changed every three days and shaken two times a day. The remaining water was subtracted from the first amount and it was checked if the rats had taken enough water or not. Starting from the day after the induction of

Table 1. Comparison of macroscopical damage scores of the groups (Mann-Whitney U test p value) (p<0.05)

Groups	Control	Ethanol	Colitis	NAC
Ethanol	0.635			
Colitis	0.001	0.001		
NAC	0.001	0.001	0.023	
Gln-NAC	0.001	0.005	0.001	0.001

Gln: Glutamine, NAC: Antioxidant N-acetylcysteine

colitis, 300 mg/kg intraperitoneal NAC was injected for 4 days. The rats continued to receive Gln together with the drinking water until sacrificed.

The rats were sacrificed on the 14th day after saline enema administration or TNBS-instillation and the distal colon was totally removed.

The Evaluation of Colonic Damage

Macroscopic Evaluation of the Colon

After the distal colon was removed under anesthesia by median laparotomy, it was opened longitudinally from the antimesenteric side of colon and the content of the colon was cleared with saline. Wallace macroscopic damage scale was used by a pathologist blinded to the content of the research.¹⁰

Histopathological Evaluation of the Colon (Table 2)

Colon tissue samples fixed in 10% formaldehyde solution were divided into 0.5 cm pieces. Then, after the standard laboratory follow-up, 5-micron sections were prepared by embedding into paraffine for histopathological examinations and sections were stained with hematoxylin-eosin. They were evaluated by using Olympus CX41 light microscope. The histopathological examinations of the tissues were performed by a pathologist blinded to sample groups using a scale that was mentioned in the previous studies.^{11,12}

Measurement of SOD

Xanthine and xanthine oxidase (XOD) were used to generate superoxide anion radicals that react with 2-(4-iodophenyl)-3-(4-nitrophenol)-5-phenyltetrazolium chloride quantitatively to form a red formazan dye. SOD inhibits the reaction by converting the superoxide radical to oxygen. The measurement of SOD enzyme activity is performed with the % of formazan dye formation at 505 nm.^{13,14}

Measurement of MDA

Under aerobic conditions, MDA reacts with thiobarbituric acid (TBA) at 95°C and pH 3.4 and forms a pink complex. This adduct is measured by spectrophotometer at 532 nm and expressed as nanomoles/milligrams.¹⁵

Statistical Analysis

Normality was checked for each continuous variable. Since the data were not distributed normally, appropriate non-parametric test was chosen. Comparisons between groups were performed using Student's t-test for normally distributed data and Mann-Whitney U test for non-normally distributed data. Bonferroni's correction was applied ($p < 0.05/n$; where n =number of comparisons) when multiple comparisons were made. The results were presented as mean \pm SD and median (minimum-maximum). Statistical analysis was performed using SPSS v 19.0 statistical package.

Results

Macroscopic Findings (Table 1)

Control Group

No change was detected in the colon macroscopically. The macroscopic damage score was 0.

Ethanol Group

The macroscopic damage score was 0. There was no statistically significant difference between the control group and the ethanol group ($p=0.635$). There was a statistically significant difference when compared with the colitis group ($p=0.001$). Statistically significant differences were found between the ethanol group and NAC group and Gln-NAC group ($p=0.001$, $p=0.001$ and $p=0.005$, respectively).

Colitis Group

The mean macroscopic damage score was 5. The score was significantly higher when compared with the control group ($p < 0.05$).

NAC Group

The mean macroscopic damage score was 3.5. A higher macroscopic damage score was found in the NAC group compared to the control group ($p < 0.05$). On the other hand, a lower macroscopic score was measured than the colitis group ($p < 0.023$). When two groups receiving Gln were compared, it was seen that damage score was higher ($p < 0.05$).

Gln-NAC Group

The mean macroscopic damage score was 4.5. Higher scores were measured compared to the control group ($p < 0.05$), whereas the scores were lower than the colitis group ($p < 0.05$). Macroscopic scores were lower compared to the NAC administered groups ($p < 0.05$).

Histopathological Findings (Table 2, Figure 1)

Control Group

The score was 0 in the histopathological evaluation of the control group.

Ethanol Group

The mean damage score was 0. There was no statistically significant difference compared to the control group ($p=0.635$). The score was significantly lower compared with the TNBS-E administered groups ($p < 0.05$ for all).

Colitis Group

The highest damage score was measured in this group (median: 6). This value was higher than all treated groups ($p < 0.05$).

Table 2. Comparison of histological damage scores of the groups (Mann-Whitney U test p value) ($p < 0.05$)

Groups	Control	Ethanol	Colitis	NAC
Ethanol	0.635			
Colitis	0.001	0.001		
NAC	0.001	0.001	0.004	
Gln-NAC	0.001	0.046	0.001	0.001

Gln: Glutamine, NAC: Antioxidant N-acetylcysteine




Figure 1. Histopathologic appearances of colon specimens: (a) Control group; normal colonic tissue HE x100, (b) Colitis group (TNBS); full-thickness colon necrosis (HE x40), (c) NAC group (TNBS + NAC); mild chronic inflammation (HE x100), (d) Gln-NAC group (TNBS + Gln-NAC); mild inflammation and edema (HE x40)

HE: Hematoxylin eosin, TNBS: Trinitrobenzene sulfonic acid, NAC: Antioxidant N-acetylcysteine, Gln: Glutamine

Distribution of values of SOD according to groups




Figure 2. Mean and 95% CI of SOD according to the study groups
SOD: Superoxide dismutase, CI: Confidence interval, NAC: Antioxidant N-acetylcysteine, Gln: Glutamine

NAC Group

When compared to with the control and ethanol groups, higher histopathological damage scores were measured (median: 2, $p < 0.05$). The score was lower than the colitis group ($p < 0.05$).

Gln-NAC Group

A slightly higher histopathological damage score was measured than the control and ethanol groups (median: 1, $p < 0.05$). The damage scores were lower in this group than the colitis and NAC groups ($p < 0.05$).

The Results of SOD Activity Measurement (Figure 2)

Ethanol Group

A statistically significant increase in SOD activity was observed in this group compared to the control group ($p = 0.007$). While there was no statistically significant difference between ethanol group and NAC group, a lower SOD activity was measured in the ethanol group compared with the TNBS-E administered groups ($p = 0.224$).

Colitis Group

The highest SOD activity was measured in this group. The SOD activity was lower in the treated groups than in the colitis group.

NAC Group

A higher SOD activity was detected compared to the control group ($p < 0.017$). The lowest SOD activity was measured in the NAC group among the treated groups.

GLN-NAC Group

The results in this group were higher than NAC group ($p = 0.001$).

The Results of MDA Measurement (Figure 3)

Ethanol Group

The values in the ethanol group were higher than in the control group ($p = 0.01$). The values were lowest when compared to all TNBS-E administered groups.

Colitis Group

The highest MDA level was obtained in this group. The values were found to be statistically higher than the treated groups.

NAC Group

Similar with SOD activity, the lowest MDA level was obtained in this group. No statistically significant difference was found between this group and ethanol group ($p = 0.201$).

Gln-NAC Group

MDA levels in this group were higher than the control group ($p = 0.001$). When compared to the colitis group, MDA levels were found to be statistically and significantly decreased ($p = 0.001$).

Distribution of values of MDA levels according to groups




Figure 3. Mean and 95% CI of MDA according to the study groups
MDA: Malondialdehyde, CI: Confidence interval, NAC: Antioxidant N-acetylcysteine, Gln: Glutamine

Discussion

Today, there is no definitive treatment for IBD except for surgery. Many clinical and experimental studies are performed to establish new treatment protocols. Oxidative stress is believed to play a key role in the pathogenesis of mucosal damage of IBD. In some of these studies, antioxidant substances have been investigated. In our study, we investigated the effect of NAC on oxidative stress and morphological changes in IBD, and the possible synergistic interaction of adding Gln, which is one of the precursors of GSH and an important antioxidant, such as cysteine in the composition of NAC.

In this study, colitis was induced with TNBS-E as shown in the literature. Expected findings were obtained in all groups with administered TNBS-E in compliance with the expected results and damage scores in all groups were higher than the control group. Unlike other studies, the efficacy of ethanol was evaluated as a given mucosal barrier breaker. The results showed that ethanol did not cause morphological damage, however, it increased SOD activity and MDA levels. Some studies evaluating the effect of ethanol on other systems have reported that altered MDA and SOD activities.^{16,17,18,19} We think that ethanol in the TNBS-induced colitis model increased the SOD activity and MDA levels and it may affect the evaluation of the results obtained.

The effect of NAC on experimental colitis model has been investigated by many researchers. Siddiqui et al.,²⁰ investigated the effect of NAC, mesalamine and NAC + mesalamine on mucosal recovery in TNBS-E-induced colitis model. It was showed that mesalamine + NAC reduced

colonic inflammation and ulceration, and provided some curative effects on mucosa compared to their individual use. Besides, it was reported that combined treatment resulted in significant myeloperoxidase activity inhibition.²⁰ In another study conducted on a group of people, NAC and mesalamine were administered to some patients, and mesalamine and placebo were administered to another group. Consequently, better clinical results were observed with decrease of chemokines in the mesalamine and NAC group.²¹ As noted in other studies, lower scores were obtained macroscopically and histopathologically in the damaged colon mucosa compared to the colitis model upon the application of NAC ($p < 0.05$). The macroscopic and histopathological results obtained by adding Gln to NAC were better.

Free oxygen radicals form lipid peroxide radicals by causing lipid peroxidation. The resulting lipid peroxide radicals play an important role in the formation of the inflammatory process. Antioxidants are protective compounds against the harmful effects of free oxygen radicals. One of the most important enzymatic antioxidants is SOD. These enzymes are part of the natural defense system. The increased MDA levels in tissues reflect the level of lipid peroxidation.²² Kurutas et al.,²³ found that NAC treatment protects the organism from the noxious substances of lipid peroxidation in acetic acid-induced colitis models. In our study, MDA levels were measured lower with NAC application compared to the colitis model. On the other hand, it was statistically significantly higher compared to the NAC group.

Two cytoplasmic enzymes, SOD and myeloperoxidase, protect the cell from oxidizing agents by breaking down superoxide anions and hydrogen peroxide. SOD reduces the oxidative stress and inflammatory response mediators.²⁴ SOD acts as an important preservative to prevent the damaging effects of superoxide anion radicals. Kuralay et al.,²⁵ showed that SOD levels increased in response to oxidative stress in the experimental colitis model and that this increase was reduced by antioxidant agents. Kruidenier et al.,²⁶ demonstrated that Cu/Zn SOD and MnSOD levels of the colonic mucosa in patients with IBD were higher than the control group. In this study, we observed lower SOD levels in the NAC group than in the colitis group. Similar with the other studies, it was concluded in our study that the antioxidant application had a positive effect on the colitis model in terms of treatment. Compared to the colitis model, a greater reduction was observed in the SOD level in the NAC + Gln group. However, contrary to expectations, the addition of Gln to NAC treatment did not provide more positive results. One reason for this is that Gln was mixed with drinking water and it might not reach enough tissue concentration.

Conclusion

In conclusion, TNBS-E-induced colitis model is easy to perform, but ethanol affects SOD activity and MDA levels. The administration of NAC appears to have beneficial effects on TNBS-E induced colitis, as indicated by decreased expression of SOD activity and MDA levels and by the scores that were used in the morphological evaluation.

Ethics

Ethics Committee Approval: The study was approved by the Çukurova University Animal Experiments Local Ethics Committee (approval number: 22.12.2016/11)

Informed Consent: Experimental animal study.

Peer-reviewed: Externally peer-reviewed.

Authorship Contributions

Surgical and Medical Practice: İ.C.E., A.R., K.D., A.G.S.,
Concept: O.A., İ.C.E., Design: İ.C.E., O.Y., S.M., Data
Collection or Processing: İ.C.E., F.D., G.S., Analysis or
Interpretation: G.S., İ.C.E., A.R., Literature Search: K.D.,
O.Y., A.G.S., İ.C.E., Writing: İ.C.E., A.R.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study received no financial support.

References

- Balmus IM, Ciobica A, Stanciu C. The implications of oxidative stress and antioxidant therapies in inflammatory bowel disease: clinical aspects and animal models. *The Saudi J Gastroenterol* 2016;22:3-17.
- Nieto N, Torres MI, Fernandez MI, Giron MD, Rios A, Suarez MD, et al. Experimental ulcerative colitis impairs antioxidant defense system in rat intestine. *Digestive diseases and sciences* 2000;45:1820-1827.
- Sido B, Hack V, Hochlehnert A, Lipps H, Herfarth C, Dröge W. Impairment of intestinal glutathione synthesis in patients with inflammatory bowel disease. *Gut* 1998;42:485-492.
- Koksal HM, Erturk AD, Celayir MF, Yıldırım S, Oner M, Baykan A. The Effects of Enoxaparine on Anastomoic Healing in Rats with Chemical Induced Inflammatory Bowel Disease. *Turk J of Colorectal* 2009;19:14-23.
- Kara E, Sakarya A, Var A, Vatansever S. The effect of low-dose methotrexate on endothelial dysfunction and programmed cell death (apoptosis) in 2,4,6 trinitrobenzenesulphonic acid (tnbsa)- induced model of experimental colitis. *Turk J Colorectal* 2010;20:66-75.
- Wengrower D, Liakim R, Karmeli F, Razin E, Rachmilewitz D. Pathogenesis of ulcerative colitis (UC): enhanced colonic formation of inositol phosphates (IP) and platelet activating factor (PAF). *Gastroenterology* 92:169-198.
- Güllüoğlu BM, Kurtel H, Güllüoğlu MG, Aktan AO, Yeğen C, Dizdaroğlu F, et al. Bombesin Ameliorates Colonic Damage in Experimental Colitis. *Dig Dis Sci* 1999;44:1531-1538.
- Gultekin FA, Emre AU, Celik SK, Barut F, Tali U, Sumer D, et al. Effects of Humanin on Experimental Colitis Induced by 2,4,6- trinitrobenzene Sulphonic Acid in Rats. *Saudi J Gastroenterol* 2017;23:105-111.
- Keshavarzian A, Kanofsky JR, Ibrahim CM, Doria MI: Excessive generation of reactive oxygen metabolites by colonic mucosa in experimental colitis: Analysis using chemiluminescence probe. *FASEB J* 1990;4:761.
- Wallace JL, MacNaughton WK, Morris GP, Beck PL. Inhibition of leukotriene synthesis markedly accelerates healing in a rat model of inflammatory bowel disease. *Gastroenterology* 1989;96:29-36.
- Morris GP, Beck PL, Herridge MS, Depew WT, Szwczuk MR, Wallace RL. Hapten-induced model of chronic inflammation and ulceration in the rat colon. *Gastroenterology* 1989;96:795-803.
- Mei Q, Yu J-P, Xu J-M, Wei W, Xiang L, Yue L. Melatonin reduces colon immunological injury in rats by regulating activity of macrophages. *Acta Pharmacologica Sinica* 2002;23:882-886.
- Fridovich I. Superoxide Radicals and Superoxide Dismutases. *Annu Rev Biochem* 1995;64:97-112.
- Lowry OH, Rosebrough NJ, Farr AL, Randall RJ. Protein Measurement with the Folin Phenol Reagent. *J Biol Chem* 1951;193:265-275.
- Ohkawa H, Ohishi N, Tagi K. Assay for lipid peroxides in animal tissues by Thiobarbituric acid reaction. *Anal Biochem* 1979;95:351-358.
- Akkuş I, Gültekin F, Aköz M, Çağlayan O, Bahçacı S, Can UG, et al. Effect of moderate alcohol intake on lipid peroxidation in plasma, erythrocyte and leukocyte and on some antioxidant enzymes. *Clin Chim Acta* 1997;266:141-147.
- Punchard NA, Senturk H, Teare JP, Thompson RP. Resistance of erythrocytes to lipid peroxidation in alcoholic patients. *Gut* 1994;35:1753-1756.
- Dudek IM, Kedziora J, Zagorski T. Effect of ethanol on human erythrocyte superoxide dismutase activity and malonyl dialdehyde concentration. *Int J Occup Med Environ Health* 1995;8:239-243.
- Armutcu F, Gurel A, Sogut S, Aksu N, Unalacak M. Levels of erythrocyte oxidant and antioxidant parameters in people with alcohol addiction. *Firat Med J* 2004;9:50-53.
- Siddiqui A, Ancha H, Tedesco D, Lightfoot S, Stewart CA, Harty RF. Antioxidant Therapy with N-Acetylcysteine Plus Mesalamine Accelerates Mucosal Healing in a Rodent Model of Colitis. *Dig Dis Sci* 2006;51:698-705.
- Guijarro LG, Mate J, Gisbert JP, Perez-Calle JL, Marín-Jimenez I, Arriaza E, et al. N-acetyl-L-cysteine combined with mesalamine in the treatment of ulcerative colitis: Randomized, placebo-controlled pilot study. *World J Gastroenterol* 2008;14:2851-2857.
- Drapper H, Hadley M. Malondialdehyde determination as index of lipid peroxidation. *Methods, Enzymol* 1990;186:421.
- Kurutas EB, Cetinkaya A, Bulbuloglu E, Kantarceken B. Effects of antioxidant therapy on leukocyte myeloperoxidase and cu/zn-superoxide dismutase and plasma malondialdehyde levels in experimental colitis. *Mediators of Inflamm* 2005;6:390-394.
- Grisham MB, Volkmer C, Tso P, Yamada T. Metabolism of trinitrobenzene sulfonic acid by the rat colon produces reactive oxygen species. *Gastroenterology* 1991;101:540-547.
- Kuralay F, Yildiz C, Ozutemiz O. Effects of trimetazidine on acetic acid-induced colitis in female Swiss rats. *J Toxicol Environ Health A* 2003;66:169-179.
- Kruidenier L, Kuiper I, van Duijn W. Differential mucosal expression of three superoxide dismutase isoforms in inflammatory bowel disease. *J Pathol* 2003;201:7-16.



Appendix Duplication Accompanied by Acute Appendicitis

Akut Apandisit Eşlik Ettiği Apendiks Duplikasyonu

© Uğur Topal¹, © Figen Doran², © Ahmet Rencüzoğulları¹

¹Çukurova University Faculty of Medicine, Department of General Surgery, Adana, Turkey

²Çukurova University Faculty of Medicine, Department of Pathology, Adana, Turkey

ABSTRACT

Acute appendicitis is the most common non-traumatic emergency surgical pathology, and duplication of the appendix is a rare congenital anomaly usually detected incidentally during laparotomy. Since Picoli first described appendix duplex in a female patient who presented with associated anomalies, few other cases have been reported. In this study, we aimed to present a case of appendix duplication detected in a 56-year-old female renal transplant patient undergoing surgery for acute appendicitis. Surgical management of double appendix is of practical importance to avoid serious medical and legal consequences.

Keywords: Double appendix, acute appendicitis, appendectomy

ÖZ

Akut apandisit travmatik olmayan acil cerrahi gerektiren en sık patoloji olup apendikse ait duplikasyon nadir bir konjenital anomalidir ve genellikle laparotomi esnasında tesadüfen saptanır. Picoli'nin eşlik eden anomalileri olan bir kadın hastada ilk apendiks duplikasyonu olgusunu bildirdikten sonra sınırlı sayıda olgu bildirilmiştir. Bu çalışmada akut apandisit nedeniyle ameliyat ettiğimiz 56 yaşında böbrek transplantlı bir kadın hastada tespit edilen apendiks duplikasyonu olgusunu sunmak amaçlanmıştır. Apendiks duplikasyonunun cerrahi yönetimi ciddi medikal ve yasal sonuçlardan kaçınmak adına pratik öneme sahiptir.

Anahtar Kelimeler: Apendiks duplikasyonu, akut apandisit, apendektomi

Introduction

The vermiform appendix develops as a conical extension from the apex of the caecal diverticulum which arises from the antimesenteric border of the proximal part of the post arterial segment of the mid gut.¹ After Picoli and colleagues reported the first case of appendix duplex in a female patient who presented with associated anomalies, limited number of cases has been reported so far.² Recently published review reported the incidence of duplicated appendix ranging from 0.004% to 0.009%.³ In this study, we aimed to present a case of an appendix duplication which was diagnosed in a 56-year-old female renal transplant patient undergoing surgery for acute appendicitis.

Case Report

A 56-year-old female patient with a history of renal transplantation from 10 years ago was admitted to the emergency room due to fever, vomiting and abdominal pain that has been going on for 2 days. On the physical examination, there was defense and rebound in the suprapubic region and subsequently migrated the right lower quadrant. Body temperature was 3.8°C and all other hemodynamic parameters were in the normal range. In the laboratory study, the white blood cells was 17.750 mm³ and the other values were in the normal range. Abdominal tomography was compatible with acute appendicitis (Figure 1). As the patient experienced kidney transplantation with



Address for Correspondence/Yazışma Adresi: Ahmet Rencüzoğulları MD,

Çukurova University Faculty of Medicine, Department of General Surgery, Adana, Turkey

Phone: +90 532 179 82 80 E-mail: rencuzdr@gmail.com ORCID ID: orcid.org/0000-0002-5993-9536

Received/Geliş Tarihi: 12.08.2018 Accepted/Kabul Tarihi: 04.09.2018




Figure 1. An inflamed bowel segment formed a mass of 2x3 cm in size which mimicked a plastrone secondary to acute appendicitis




Figure 2. Macroscopic appearance of inflamed appendix (a), normal appendix (b) and pathologic specimen (c)




Figure 3. Microscopic view showing (H&E, X100) mucosal erosion and inflammation indicating acute appendicitis (a) and appendix with lymphoid follicles (b)

a right-oblique lumbar incision extending right lower quadrant, open surgical approach is preferred. During exploration, Type B2 appendix duplication, based on the Cave-Wallbridge classification⁵, was detected (Figure 2). The patient was examined for congenital malformation of other intra-abdominal organs, and no other malformation was identified. While one of the appendixes had the findings of acute inflammation, the other one had a normal appearance (Figure 3a and 3b). Appendectomy for both appendixes was performed. Histopathological examination of the surgical specimen confirmed the intraoperative findings of appendiceal duplication as well as findings of acute inflammation of the second appendix. On pathological examination, the appendixes measured 7x4x3.5cm and 8x3x1cm. One of the appendixes was acute perforated appendicitis, periappendicitis and the other was appendix vermiformis. Microscopic images of both appendixes are shown in the Figure 1c. The patient had a full recovery without postoperative complications and was discharged on the postoperative day 4.

Discussion

Although the normal embryogenesis of the appendix is known, the exact cause of appendiceal duplication has not been fully explained. In an attempt to explain the pathogenesis of duplication, Cave put forward two theories; (i) the persistence of a transient embryological structure and (ii) incidental appendiceal duplicity to a more general affection of the primitive midgut.⁴

Since preoperative radiological identification of duplication of appendix is difficult, computed tomography may be the best mode of imaging to identify a duplex appendix.³ The surgeons intraoperative attention and awareness are vital in terms of diagnosis. Macroscopic view and histopathological examination are substantially pathognomonic. The majority

Table 1. Cave-Wallbridge classification

Type A	Single caecum with one appendix exhibiting partial duplication.
Type B	Single caecum with two obviously separate appendixes.
B1	The two appendixes arise on either side of the ileocaecal valve in a B bird-like manner.
B2	In addition to a normal appendix arising from the caecum at the usual site, there is also a second, usually rudimentary, appendix arising from the caecum along the lines of the taenia at a varying distance from the first.
B3	The second appendix is located along the taenia of the hepatic flexure of the colon.
B4	The location of the second appendix is along the taenia of the splenic flexure of the colon.
Type C	Double caecum, each bearing its own appendix and associated with multiple duplication anomalies of the intestinal tract as well as the urinary tract.

of duplicated appendixes is believed to be silent and only discovered when one of them becomes inflamed.

In 1936, Cave classified appendiceal duplication for the first time which was modified by Wallbridge in 1963 into three types; as shown in Table 1.^{4,5}

Appendix duplications are usually detected incidentally during surgery for acute appendicitis and it was identified in the same way in our case. As performed in our case, appendectomy for both appendixes is necessary for the correct evaluation of the clinical problems of patients with subsequent abdominal pain. We noticed that our patient had type B2 appendiceal duplication based on the Cave-Wallbridge classification. Most authors agree that Type B2 duplication may be the remnant of a “transient appendix” which appears during the sixth and seventh week of embryological development, the presence of which was first reported by Kelly and Hurdon.^{3,6} Although appendiceal duplication may be associated with other congenital abnormalities⁴ any other anomalies were absent in our patient.

Conclusion

In conclusion, intraoperative detection of appendicular duplication is of great practical importance, as overlooking of this clinical scenario may lead to medical and legal issues. Surgeons, especially junior surgical residents, should be aware of the potential anatomical anomalies and malpositions of the vermiform appendix; careful inspection of the caecum should be performed to avoid potentially serious clinical and medicolegal consequences. Moreover, acute appendicitis should be considered as a part

of differential diagnosis of acute right lower quadrant pain despite history of appendectomy.

Ethics

Informed Consent: Obtained from the patient that was mentioned in the manuscript.

Peer-review: Externally peer-reviewed.

Author Contributions

Surgical and Medical Practices: U.T., A.R., Concept: U.T., F.D., A.R., Design: U.T., F.D., A.R., Data Collection or Processing: U.T., F.D., A.R., Analysis or Interpretation: U.T., F.D., A.R., Literature Search: U.T., F.D., A.R., Writing: U.T., F.D., A.R.

Conflict of Interest: The authors have no conflicts of interest including relevant financial interests, activities, relationships, and affiliations.

Financial Disclosure: The authors declared that this study received no financial support.

References

1. Moore KL. Saunders. The developing human: clinically oriented embryology with student consult online access; 9th edition. 2011.
2. Khanna AK, Appendix Vermiformis Duplex. *Postgrad Med J* 1983;59:69-70.
3. Nageswaran H, Khan U, Hill F, Maw A. Appendiceal duplication: a comprehensive review of published cases and clinical recommendations. *World J Surg* 2018;42:574-558.
4. Cave AJ. Appendix vermiformis duplex. *J Anat* 1936;70:283-292.
5. Wallbridge PH. Double appendix. *Br J Surg* 1963;50:346-347. doi:10.1002/bjs.18005022124.
6. Kelly HA, Hurdon E. Vermiform appendix and its diseases. Philadelphia; WB Saunders Co. 1905.



Idiopathic Nonobstructive Colo-colonic Intussusception: A Rare Clinical Presentation

İdiyopatik Nonobstrüktif Kolo-kolonik İntussusepsiyon: Nadir Bir Klinik Tablo

© Fatma Ayça Gültekin¹, © Ramazan Kozan¹, © Yücel Üstündağ²

¹Zonguldak Bülent Ecevit University Faculty of Medicine, Department of General Surgery, Zonguldak, Turkey

²Zonguldak Bülent Ecevit University Faculty of Medicine, Department of Internal Diseases, Division of Gastroenterology, Zonguldak, Turkey

ABSTRACT

Adult intussusception is a rare clinical condition. There is generally an etiological factor, mainly malignancy. However, the number of idiopathic cases is very low. Patients may present with obstruction or acute abdomen, but the symptoms and signs may also be nonspecific. The diagnostic process is more difficult in patients with non-specific clinical presentation. In this article, the diagnosis and treatment options of an adult idiopathic colo-colonic intussusception are discussed.

Keywords: Intussusception, adult, idiopathic, colonic diseases, laparoscopy

ÖZ

İntussusepsiyon erişkinlerde nadir görülen bir durumdur. Başta malignite olmak üzere etiyolojik faktör sıklıkla tespit edilebilmektedir. Buna karşın idiyopatik olgu sayısı oldukça azdır. Klinik tablo, intestinal obstrüksiyon veya akut karının yanı sıra spesifik olmayan semptom ve bulgular şeklinde de ortaya çıkabilmektedir. Bu hastalarda tanısal süreç daha güçtür. Etkin preoperatif değerlendirme seçilecek cerrahi tedavi şekli ve genişliğini belirlemesi açısından önem taşır. Bu yazıda yetişkin bir idiyopatik kolo-kolonik intussusepsiyon olgusu eşliğinde tanı ve tedavi seçenekleri tartışılmıştır.

Anahtar Kelimeler: İntussusepsiyon, erişkin, idiyopatik, kolonik hastalıklar, laparoskopisi

Introduction

Intussusception is most common in infants and children, but rare in adults, representing 5% of all bowel intussusceptions and 1% of all bowel obstructions.^{1,2,3,4} The large bowel is the most common site for adult intussusception, and 80% of cases are colo-colonic intussusception.⁵ A causal lesion is identified in 90% of adult intussusceptions and 44% of these lesions are malignant.⁶ However, there are few cases of adult colo-colonic intussusception without any identifiable lesion in the literature. In this article, we report a case of adult colo-colonic intussusception without any pathologic underlying cause who was treated by laparoscopic resection.

Case Report

A 41-year-old male patient was admitted to emergency department with a complaint of abdominal colic pain predominantly in the left flank for 10 days. Two day before the admission, the patient reported worsening abdominal pain, nausea and vomiting. His bowel movement was normal. He had no comorbid disease, previous abdominal surgery or family history of colorectal cancer. Physical examination revealed painful abdominal distention without signs of peritoneal irritation. Bowel sounds were present. Initial laboratory values were within normal limits. Abdominal X-rays were taken and did not show any signs of obstruction or perforation.



Address for Correspondence/Yazışma Adresi: Ramazan Kozan MD,
Zonguldak Bülent Ecevit University Faculty of Medicine, Department of General Surgery, Zonguldak, Turkey
Phone: +90 532 763 16 97 E-mail: dr.kozan@hotmail.com ORCID ID: orcid.org/0000-0002-3835-8759
Received/Geliş Tarihi: 23.09.2018 Accepted/Kabul Tarihi: 28.10.2018

Abdominal and pelvic computed tomography (CT) demonstrated thickening of the sigmoid colon wall and mild distension of the ascending and transverse colon. Magnetic resonance imaging (MRI) was performed in accordance with the recommendations of radiology department on the basis of suspicion of intussusception in CT and thickening of the wall. Abdominal and pelvic MRI revealed “bowel within bowel” colo-colonic intussusception at the level of descending colon and sigmoid colon (Figure 1). The length of the affected colon segment was about 10 cm. In the absence of signs suggestive of acute intestinal obstruction, the patient was hospitalized for further investigation. On the second day of hospitalization, colonoscopy was performed in order to exclude malignancy. In the colonoscopic examination, congested colonic mucosa completely filled the colon lumen without evidence of malignancy, but permitted the advance of the colonoscopy (Figure 2). According to the findings, the patient was prepared for laparoscopic sigmoid resection with bowel cleansing. Intraoperatively, a round mass-like appearance was observed in the sigmoid colon as a result




Figure 1. Magnetic resonance appearance of colo-colonic intussusception at the level of descending colon and sigmoid colon




Figure 2. Colonoscopic view of colo-colonic intussusception

of an invagination of the descending colon into the sigmoid colon (Figure 3). Loose adhesions were also observed between the affected bowel segment and pelvic walls (Figure 4). No other abnormality was found. Laparoscopic resection of descending colon and primary anastomosis were performed. Gross examination of the resected specimen revealed a 15 cm colo-colonic intussusception (Figure 5). Histological examination revealed non-specific mucosal inflammation and fibrotic thickening at serosal surfaces




Figure 3. A round mass like appearance in the sigmoid colon (arrow) due to colo-colonic intussusception




Figure 4. Loose adhesions between the affected bowel segment and pelvic walls



Figure 5. A macroscopic view of the resected specimen

of the colon segments. No neoplastic or benign lesion was identified. The postoperative period was uneventful and the patient was discharged on the fifth postoperative day. Written informed consent was obtained from the patient for the use of medical data.

Discussion

Adult intussusception is a rare clinic entity. The signs and symptoms of adult colo-colonic intussusception are nonspecific, and most patients present with chronic or intermittent abdominal pain, partial intestinal obstruction and gastrointestinal bleeding.⁷ In the literature, 75% of patients presented with obstruction symptoms, 5% with acute abdomen, and one third of patients had a palpable abdominal mass. In adult patients, an underlying cause is found in 75-90% of patients, and malignancy is the most common etiology. Adenocarcinoma was particularly common in adult colo-colonic intussusception.⁸

An accurate preoperative diagnosis of adult colo-colonic intussusception is relatively difficult. There is no specific laboratory finding for adult intussusception. Due to being cost effective, ultrasonography (US) is considered a first-level diagnostic technique. The characteristic ultrasonographic findings of intussusception include the “target” or “doughnut” signs in the transverse view and the “pseudo-kidney” sign or “hay-fork” sign in the longitudinal view.^{9,10} However, accuracy of US depends on status of patient body (obese patient etc.) and experience of radiologist. CT is the most valuable imaging technique to confirm a preoperative diagnosis of intussusception. The diagnostic accuracy of CT reported for intussusception is 58-100%.⁶ CT can also demonstrate any mass associated with intussusception and may help assess the location and character of the mass and its association with surrounding tissues. CT can provide diagnostic findings of intussusception including an inhomogeneous “target” or “sausage” shaped soft tissue mass with a layering effect and/or an apparent mass lesion.^{11,12} Endoscopic investigations such as colonoscopy and sigmoidoscopy can help identify both intussusception and the underlying causes (leading point) of intussusception and collect a histology sample.⁶ However, in case of intestinal obstruction, endoscopy is not advisable due to the high risk of perforation. As the patient in our case did not show typical symptoms associated with intestinal obstruction, we performed colonoscopy.

The treatment of choice for adult intussusception is surgical resection of the affected bowel segment. The extent of resection depends on the underlying cause of intussusception, and malignant lesions require oncologic resection.⁶ In patients with right sided intussusception (ileo-

colonic, ileo-cecal and colo-colonic), construction with primary anastomosis is recommended, whereas Hartman procedure is more appropriate surgical option for left-sided or rectosigmoid intussusceptions, especially in the emergency setting.⁷ In contrast to most pediatric patients, preoperative reduction of intussusception is not feasible for adult patients, and carries increased risk of perforation, tumor seeding (particularly intussusception associated with malignant lesions) and anastomotic complications. However, if preoperative diagnosis of lesion is benign and the bowel wall looks viable and healthy, reduction may be performed in order to limit extent of bowel resection.

Successful laparoscopic resections in adult intussusception cases have been reported in the literature.^{13,14,15} Laparoscopy can be used for both diagnostic and operative purposes. Once the diagnosis of intussusception and underlying disease is confirmed laparoscopically, the surgeon can perform laparoscopic resection. However, the characteristics of intussusception (location and extent), underlying cause and the expertise of the surgeon affects the outcome of laparoscopic surgery. In this case, preoperative diagnosis of intussusception was performed by CT, MRI and colonoscopy, and laparoscopic resection was performed successfully.

Conclusion

In conclusion, idiopathic colo-colonic intussusception without intestinal obstruction is a rare clinical phenomenon. Non-specific clinical presentation makes the preoperative diagnosis of intussusception difficult. However, CT can help to establish preoperative diagnosis. Resection of the affected bowel segment is recommended and choice of surgical technique (laparoscopic or open) should be made based on characteristics of intussusception and the experience of the surgeon.

Ethics

Informed Consent: Written informed consent was obtained from patient in this case.

Peer-review: External and internal peer-reviewed.

Authorship Contributions

Surgical and Medical Practices: F.A.G., R.K., Y.U., Concept: F.A.G., Y.U., Design: F.A.G., R.K., Data Collection or Processing: F.A.G., Y.U., Analysis or Interpretation: F.A.G., R.K., Literature Search: F.A.G., R.K., Writing: F.A.G., R.K.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study received no financial support.

References

1. Rutherford CL, Alkhaffaf B, Massa E, Turner P. Colo-colic intussusception secondary to lipomatous polyp in an adult. *BMJ Case Rep* 2013;2013:bcr2012008037.
2. El-Sergany A, Darwish A, Mehta P, Mahmoud A. Community teaching hospital surgical experience with adult intussusception: Study of nine cases and literature review. *Int J Surg Case Rep* 2015;12:26-30.
3. Honjo H, Mike M, Kusanagi H, Kano N. Adult intussusception: a retrospective review. *World J Surg* 2015;39:134-138.
4. Lianos G, Xeropotamos N, Bali C, Baltogiannis G, Ignatiadou E. Adult bowel intussusception: presentation, location, etiology, diagnosis and treatment. *G Chir* 2013;34:280-283.
5. Eisen LK, Cunningham JD, Aufses AH Jr. Intussusception in adults: institutional review. *J Am Coll Surg* 1999;188:390-395.
6. Marinis A, Yiallourou A, Samanides L, Dafnios N, Anastasopoulos G, Vassiliou I, et al. Intussusception of the bowel in adults: a review. *World J Gastroenterol* 2009;15:407-411.
7. Azar T, Berger DL. Adult intussusception. *Ann Surg* 1997;226:134-138.
8. Begos DG, Sandor A, Modlin IM. The diagnosis and management of adult intussusception. *Am J Surg* 1997;173:88-94.
9. Boyle MJ, Arkell LJ, Williams JT. Ultrasonic diagnosis of adult intussusception. *Am J Gastroenterol* 1993;88:617-618.
10. Weissberg DL, Scheible W, Leopold GR. Ultrasonographic appearance of adult intussusception. *Radiology* 1977;124:791-792.
11. Gayer G, Apter S, Hofmann C, Nass S, Amitai M, Zissin R, et al. Intussusception in adults: CT diagnosis. *Clin Radiol* 1998;53:53-57.
12. Gayer G, Zissin R, Apter S, Papa M, Hertz M. Pictorial review: adult intussusception-a CT diagnosis. *Br J Radiol* 2002;75:185-190.
13. Ishibashi Y, Yamamoto S, Yamada Y, Fujita S, Akasu T, Moriya Y. Laparoscopic resection for malignant lymphoma of the ileum causing ileocecal intussusception. *Surg Laparosc Endosc Percutan Tech* 2007;17:444-446.
14. Akatsu T, Niihara M, Kojima K, Kitajima M, Kitagawa Y, Murai S. Adult colonic intussusception caused by cecum adenoma: successful treatment by emergency laparoscopy: report of a case. *Surg Today* 2007;37:694-697.
15. Palanivelu C, Rangarajan M, Senthilkumar R, Madankumar MV. Minimal access surgery for adult intussusception with subacute intestinal obstruction: a single center's decade-long experience. *Surg Laparosc Endosc Percutan Tech* 2007;17:487-491.



Penetrating Sharp Object Injury in the Gluteal Region and Small Bowel Perforation Due to Pelvic Penetration: A Case Report

Gluteal Bölgede Kesici Delici Alet Yaralanması: Pelvis Nafiziyeti Sonucu İnce Barsak Perforasyonu: Olgu Sunumu

© Ergün Yüksel¹, © Mehmet Akif Üstüner², © Ahmet Karayiğit¹, © Lütfi Doğan¹

¹University of Health Sciences, Dr. Abdurrahman Yurtaslan Ankara Training and Research Hospital, Department of General Surgery, Ankara, Turkey

²Türkiye Yüksek İhtisas Training and Research Hospital, Clinic of Gastroenterology Surgery, Ankara, Turkey

ABSTRACT

Penetrating sharp object injuries in the gluteal region are common. However, it is very rare that these injuries lead to perforation of the small bowel by pelvic penetration, which is generally overlooked. As penetration depth increases, there may be life-threatening injuries such as vascular injury, bowel injury and genitourinary injury depending on the localization of injury site. The case report addressed in the present research focuses on a patient who was admitted to the hospital because of a penetrating sharp object injury in the gluteal region.

Keywords: Penetrating sharp object injuries, gluteal region, perforation of the small intestine

ÖZ

Gluteal bölgedeki kesici delici alet yaralanmaları sık görülmektedir. Ancak bu yaralanmaların pelvise penetrasyon göstererek ince barsak perforasyonuna sebep olması çok nadirdir. Penetrasyon derinliği arttıkça, yaralanma yerinin lokalizasyonuna bağlı olarak vasküler yaralanma, barsak yaralanması, genitoüriner yaralanma gibi hayatı tehdit eden yaralanmalar olabilir. Bu olgu sunumunda hastaneye gluteal bölgeye kesici delici alet yaralanmasıyla başvuran olgumuzu sunduk.

Anahtar Kelimeler: Kesici delici alet yaralanması, gluteal bölge, ince barsak perforasyonu

Introduction

The gluteal region is anatomically located between iliac wings, inferior gluteal folds and bilateral greater trochanter.¹ Penetrating sharp object injuries (PSOIs) in the gluteal region are rare and constitute 2-3% of all injuries.² These injuries generally do not pose serious health problems. However, visceral and vascular injuries leading to life-threatening events can be observed in one-quarter of the cases.³ To the best of our knowledge, little research has been conducted on visceral and vascular injuries in the gluteal region. In this study, we present a case of small bowel perforation arising from an injury in the gluteal region. We hope our study

will bring more interest to the subject and contribute to the scarce literature.

Case Report

A 24-year-old male patient was admitted to the Emergency Department of Dr. Abdurrahman Yurtaslan Ankara Oncology Training and Research Hospital with a PSOI in the gluteal region. Physical examination of the patient revealed 1 cm incision in the superior zone of the left gluteal region at 10 o'clock and eviscerated gangrenous ileum loop in the superior zone of the right gluteal region at 9 o'clock (Figure 1). The vitals were stable. A urinary bladder catheter was inserted and



Address for Correspondence/Yazışma Adresi: Mehmet Akif Üstüner MD,

Türkiye Yüksek İhtisas Training and Research Hospital, Clinic of Gastroenterology Surgery, Ankara, Turkey

Phone: +90 507 924 62 87 E-mail: dr_ustuner@hotmail.com ORCID ID: orcid.org/

Received/Geliş Tarihi: 30.10.2018 Accepted/Kabul Tarihi: 04.01.2019

no macroscopic hemorrhage was observed. Lower extremity peripheral pulses were normal by palpation on both sides. There was no motor deficit in both lower extremities. No abnormality was found in the digital rectal examination.

Abdominal examination revealed muscular guarding and rebound tenderness. Abdominal examination findings and gangrenous prolapse of the ileal segment mandated surgical intervention. Laparotomy was performed via a midline abdominal infraumbilical incision. At exploration,




Figure 1. A 1 cm penetrating injury in the superior zone of the left gluteal region at 10 o'clock; eviscerated gangrenous ileum loop in the superior zone of the right gluteal region at 9 o'clock




Figure 2. The intestinal loop with 3-4 perforations which was incarcerated through obturator canal which was opened after injury



Figure 3. Postoperative gluteal injury sites

approximately 200 cc hemorrhagic fluid was present in the abdomen. It was observed that approximately 15 cm incarcerated small intestinal loop, 20 cm proximal to the ileocaecal valve, had slipped out through the penetrated obturator canal during the injury. The incarcerated intestinal loop was taken back into the abdominal cavity, and three perforated areas and necrosis were observed (Figure 2). The injured segment was resected and end-to-end anastomosis was performed.

In further exploration, no injury was detected to the iliac, femoral and obturator arteries or veins, aorta and vena cava inferior. The urinary bladder and ureter were intact. After rectal tube placement, it was concluded that the rectum was not injured. The abdominal cavity was irrigated with copious isotonic fluid. The peritoneum in the injured area was debrided and repaired. A drain was placed in the right pararectal space. The injured areas in the gluteal region had already been explored preoperatively, and the operation was completed by placing a Penrose drain in both injury sites (Figure 3).

On postoperative day 2, the possibility of a rectal injury was re-evaluated by rectoscopy. No injury was detected. Neurosurgery consultation recommended electromyography to rule out possible sciatic nerve damage and no deficit was detected by EMG. The bowel movements started on postoperative day 2 and he started oral feeding on day 3. At the end of postoperative day 6, he was discharged by surgical healing.

Discussion

PSOIs are as old as the history of humanity. Nowadays, PSOIs occur with the use of different objects in our country and around the world. According to the study conducted by Üstüner et al.,⁴ PSOIs are usually performed by young men (16-35 years old), most often in summer, during may and between 12.00-04.00 a.m. Among these injuries, gluteal injuries are the third most common injuries following thoracic and abdominal injuries.⁵ Depending on the type of injury, patients are generally discharged with antibiotics after closure of the wound with primary suture. However, there may be vascular, intestinal, genitourinary and sciatic nerve injuries depending on the depth and trajectory of penetration.⁶ The upper zone and the lower zone are formed when the gluteal region is divided into two from the intertrochanteric line. Mortality and morbidity of the upper zone injuries are higher than the lower zone.³ Angiography and surgical intervention may be required for diagnosis and treatment.⁷ In our case, the injury occurred in the upper zone and the patient was operated.

There is no consensus on the treatment algorithm of penetrating injuries to the gluteal region and the

management is still to be evolved. Treatment begins with ABCDE. First, CT is performed to investigate the presence of intraperitoneal fluid in patients with shock. If positive, laparotomy is performed and the localization of the bleeding is determined if negative. Packing and angioembolization are performed depending on the bleeding site, and surgery is performed if the bleeding could not be stopped. In patients without shock, CT is also the initial imaging method. Surgery, angioembolization or observation could be preferred according to the observed bleeding and abdominal injuries.²

In 2014, Lunevicius et al.,⁸ updated the algorithm and recommended evaluation in the following order: general inspection of the patient, controlling the femoral pulse, neurological examination of the lower extremities, examination of urethra and rectum, and then rectoscopy was suggested after the patient became stable.⁹ Radiologists recommend contrast-enhanced rectal CT if there is still clinical suspicion of rectal injury despite rectoscopy.⁶ There is no standard treatment for pelvic wounds.¹⁰

In our case, the examination of the rectum, genital organs, femoral and distal pulses were performed. Since there was an incarcerated intestinal loop seen outside of the gluteal region and findings of acute abdomen, the surgery was performed. On the postoperative day 2, rectum was checked using rectoscopy. In PSOIs, the mortality rate is 0-2% in the posterior abdomen region, 0-4.4% in the anterior abdomen region, 2.1% in the thoraco-abdominal region, and 2.5-5.6% in the thorax region. The mortality rate in gluteal injuries is 2.9%, which is quite high compared to other parts of the body.²

Conclusion

In conclusion, PSOIs in the gluteal region may lead to life-threatening damages. Small bowel perforation is one of these damages and should be kept in mind though it is rarely observed.

Ethics

Informed Consent: Informed consent was obtained from the patient.

Peer-review: Externally peer-reviewed.

Authorship Contributions

Surgical and Medical Practices: E.Y., A.K., Concept: M.A.Ü., Design: M.A.Ü., L.D., Data Collection or Processing: A.K., Analysis or Interpretation: M.A.Ü., L.D., Literature Search: E.Y., Writing: M.A.Ü.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study received no financial support.

References

1. Ivatury RR, Rao PM, Nallathambi M, Gaudino J, Stahl WM. Penetrating gluteal injuries. *J Trauma* 1982;22:706-709.
2. Lunevicius R, Schulte KM. Analytical review of 664 cases of penetrating buttock trauma. *World J Emerg Surg* 2011;6:33.
3. Mercer DW, Buckman RF Jr, Sood R, Kerr TM, Gelman J. Anatomic considerations in penetrating gluteal wounds. *Arch Surg* 1992;127:407-410.
4. Üstüner MA, İlhan E, Yıldırım M, Aykas A, Şenlikci A, Degerli V, et al. Penetrating stab wounds to the abdominal wall: retrospective analysis of 131 cases: can the number of unnecessary laparotomies be reduced? *Tepecik Egit Hast Derg* 2015;25:143-150.
5. Susmallian S, Ezri T, Elis M, Dayan K, Charuzi I, Muggia-Sullam M. Gluteal stab wound is a frequent and potentially dangerous injury. *Injury* 2005;36:148-150.
6. Campion T, Cross S. The spectrum of injuries in buttock stab wounds. *Clinical Radiology* 2017;543-551.
7. Makrin V, Sorene ED, Soffer D, Weinbroum A, Oron D, Kluger Y. Stab wounds to the gluteal region: a management strategy. *J Trauma* 2001;50:707-710.
8. Lunevicius R, Lewis D, Ward RG, Chang A, Samalavicius NE, Schulte KM. Penetrating injury to the buttock: an update. *Techn Coloproctol* 2014;18:981-992.
9. Keğin M, Alçiçek S, Er M, Yıldırım E, Uçar N. Ileum Perforation after a Single Gluteal Stab Wound. *Jarem* 2015;5:125-127.
10. Uğur M, Oruç C, Yıldız İ, Koca YS, Daban U. The role of co-administration of damage control surgery and vacuum-assisted closure in the treatment of perineal wounds. *Turk J Surg* 2018;34:229-230.



Midgut Nonrotation in an Adult Patient and Ladd's Procedure

Yetişkin Hastada Midgut Nonrotasyonu ve Ladd Prosedürü

© Huriye Hande Aydın¹, © Özgen Işık², © Tuncay Yılmazlar²

¹New York University, School of Medicine, Department of Surgery, Division of General Surgery, New York, United States

²Uludağ University Faculty of Medicine, Department of Surgery, Bursa, Turkey

ABSTRACT

Midgut malrotation is a congenital anomaly that usually occurs during infancy. The most common type is the "classic" form, which is defined as complete failure of rotation of both proximal (duodenojejunal limb) and distal (cecocolic) small bowel loops (nonrotation). Delayed diagnosis of midgut malrotation anomalies may lead to progression to catastrophic complications including volvulus and ischemic necrosis. Ladd's procedure is the standard of care. We aimed to report a brief case of a midgut nonrotation in an adult male patient.

Keywords: Midgut, malrotation, anomaly

ÖZ

Midgut malrotasyonu çoğunlukla doğum sonrası erken dönemde görülen konjenital bir anomalidir. En sık tipi hem proksimal (duodenojejunal bacak) hem de distal (çekokolik) ince barsak anslarının rotasyonunun olmaması ile karakterize olan klasik tiptir. Midgut malrotasyon anomalilerinde gecikmiş tanı volvulus ve iskemik nekrozu da içerebilen katastrofik komplikasyonlara yol açabilir. Ladd prosedürü standart tedavi şeklindedir. Burada yetişkin bir erkek hastada midgut nonrotasyon olgusunu bildirmeyi amaçladık.

Anahtar Kelimeler: Midgut, malrotasyon, anomaly

Introduction

Midgut malrotation is a congenital anomaly and usually occurs in infancy. The most common type is the "classic" form of malrotation defined as complete failure of rotation of both proximal (duodenojejunal limb) and distal (cecocolic) small bowel loops (nonrotation).¹ Although this pathology may be a common diagnosis for pediatric surgeons, this may not be the case for surgeons treating adults in general. Delayed diagnosis of midgut malrotation anomalies can cause progression to catastrophic complications including volvulus and ischemic necrosis. We aimed to report a case of midgut nonrotation in an adult male patient and to increase familiarity with this disease among surgeons treating adults.

Case Report

A 29 year-old male patient was referred to our clinic from an outside hospital with a suspected inflammatory bowel disease due to ongoing abdominal pain, nausea, vomiting and constipation for the last 5 days. He was otherwise healthy with a past medical history significant only for benign prostatic hypertrophy and no prior abdominal surgeries. He was admitted to our clinic with an initial diagnosis of intestinal obstruction.

On physical examination, he was afebrile and hemodynamically stable with no signs of acute abdomen. Computerized tomography scan of the abdomen showed findings consistent with proximal small bowel obstruction. His initial management consisted of intravenous (IV)



Presented in: This study was presented as a poster at the XVI. Turkish National Colon and Rectal Surgery Congress in Antalya, Turkey in May 15-20, 2017.

Address for Correspondence/Yazışma Adresi: Özgen Işık MD,

Uludağ University Faculty of Medicine, Department of Surgery, Bursa, Turkey

Phone: +90 535 895 23 05 E-mail: ozgen006@yahoo.com ORCID ID: orcid.org/ 0000-0002-9541-5035

Received/Geliş Tarihi: 30.12.2018 Accepted/Kabul Tarihi: 12.01.2019



Figure 1. Intraoperative view of small bowel and colon (after excision of Ladd bands and appendectomy)

fluids, IV antibiotics and nasogastric decompression. Over a 24-hour period, the nasogastric tube output was 2 liters and clinical findings of mechanical obstruction did not improve. He consented for an exploratory laparotomy in the emergency settings. Intraoperatively, small bowel was found to be segregated into the right abdomen, consistent with midgut nonrotation anomaly. Subsequently, following steps were performed: counterclockwise detorsion of the small bowel, internal hernia reduction and take down of Ladd's band in the right upper quadrant (RUQ). Following these steps, cecum and appendix were identified in the left lower quadrant, and an appendectomy was also performed. Relocation of the duodenum to the RUQ was undertaken and subsequent to that colon was placed in the left abdomen. The affected small bowel segment was re-examined in terms of perfusion, and resection was deemed not necessary per operating surgeon (Figure 1). The patient's abdomen was closed primarily and he was transferred to the surgical floor. Postoperative course was uneventful and patient was discharged home on postoperative day 7.

Discussion

Intestinal malrotation occurs due to the lack of normal rotation and fixation of the embryologic gut during the first trimester of gestation.² Most of the patients with intestinal malrotation present during infancy with acute clinical presentation of intestinal obstruction. Although rare, adult patients more commonly present with chronic symptoms including intermittent vomiting, abdominal pain, weight loss, and food intolerance that lasts for more than 6 months.³ High index of suspicion is required to make an early diagnosis and intervention and subsequently to prevent progression to bowel necrosis with midgut volvulus in adult cohort. Imaging studies are usually suggestive of the disease, but in most cases it is diagnosed intraoperatively. An upper

gastrointestinal series can be used to establish the diagnosis preoperatively; nevertheless surgery shouldn't be postponed in patients with acute clinical presentation for further imaging. The choice of treatment depends on the severity of the presentation. Ladd's procedure is the standard of care and it can be scheduled electively in patients with chronic disease or as an emergency in the acute setting.² Nasogastric decompression, volume resuscitation and antibiotics are important aspects of preoperative preparation. Ladd's procedure is divided into the following 5 steps: initially, the mesentery should be assessed for any twists and signs of ischemia, and counterclockwise detorsion of volvulus is achieved if applicable. Following this, Ladd bands, which are fibrous bands between cecum and duodenum that can cause duodenal obstruction, should be divided. In addition, intermesenteric fibrous bands between bowel loops should be identified by running the entire bowel and should be divided. Due to higher incidence of aberrant location of the appendix, appendicitis in patients with malrotation was found to have a higher complication rate. An appendectomy is routinely performed as a part of the Ladd's procedure. The final step consists of placing the bowel in the corrected anatomic position as small bowel in the right abdomen, colon in the left abdomen, and cecum in the left upper quadrant.² Postoperative morbidity and reoperation rates are high in adult patients when compared to pediatric cohort. This might be due to longer time to diagnosis and more advanced disease.⁴

We shared an acute adult presentation of an intestinal malrotation with midgut volvulus. Surgeons primarily serving adult patient cohorts should keep malrotation as a differential diagnosis. Intestinal malrotation should be suspected in patients with abdominal pain with or without accompanying bowel obstruction symptoms. In addition, a more subtle presentation with chronic nonspecific gastrointestinal symptoms may be related to intestinal malrotation.

Ethics

Informed Consent: Written informed consent obtained

Peer-review: Externally peer-reviewed.

Authorship Contributions

Surgical and Medical Practices: Ö.I., T.Y., Concept: Ö.I., T.Y., Design: Ö.I., T.Y., Data Collection or Processing: Ö.I., T.Y., Analysis or Interpretation: Ö.I., H.H.A., Literature Search: Ö.I., H.H.A., Writing: Ö.I., H.H.A.

Conflict of Interest: No conflict of interest was declared by the authors.

Financial Disclosure: The authors declared that this study received no financial support.

References

1. Dietz DW, Walsh RM, Grundfest-Broniatowski S, Lavery IC, Fazio VW, Vogt DP. Intestinal malrotation: a rare but important cause of bowel obstruction in adults. *Dis Colon Rectum* 2002;45:1381-1386.
2. Buchmiller T. Intestinal malrotation in adults 2017 [Available from: https://www-uptodate-com.ezproxy.med.nyu.edu/contents/intestinal-malrotation-in-adults?search=intestinal%20malrotation&source=search_result&selectedTitle=2~41&usage_type=default&display_rank=2#H10198704]
3. Yanez R, Spitz L. Intestinal malrotation presenting outside the neonatal period. *Arch Dis Child* 1986;61:682-685.
4. Durkin ET, Lund DP, Shaaban AF, et al. Age-related differences in diagnosis and morbidity of intestinal malrotation. *J Am Coll Surg* 2008;206:658-663.

NOTES

A series of horizontal dotted lines for writing notes, spanning the width of the page below the 'NOTES' header.